

**MEASUREMENT OF THE $Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ JET PRODUCTION CROSS
SECTIONS IN $p\bar{p}$ COLLISIONS AT $\sqrt{s} = 1.96$ TeV
DRAFT v1.9**

BY

MARC BUEHLER

B.S. (University of Basel, Switzerland) 1992
M.S. (University of Heidelberg, Germany) 1997

THESIS

Submitted in partial fulfillment of the requirements
for the degree of Doctor of Philosophy in Physics
in the Graduate College of the
University of Illinois at Chicago, 2005

Chicago, Illinois

Copyright by

Marc Buehler

2005

To Dasha

ACKNOWLEDGMENTS

First, I wish to thank my thesis advisor, Nikos Varelas, who guided and supported me throughout the years. Thanks to the other members of the UIC group at DØ - Mark Adams, Cecilia Gerber, and Elizaveta Shabalina - for their assistance. Special thanks to Alan Stone for his help.

I am also indebted to many people in the Higgs Group at DØ ; to the Higgs group conveners Gregorio Bernardi, and Avto Kharchilava; to Kazu Hanagaki, who taught me to never surrender; to Jodi Wittlin, who tried to convince me that Higgs bosons are red. I would especially like to thank Suyong Choi for his help and advice.

I was lucky to work on the trigger system alongside Adam, Burair, Josh K., Miroslav, and Terry T.

Of course, this work would not have been possible without the help and support from the hard working DØ Collaboration (see Appendix A).

On a more personal note, I would like to thank the friends who kept me sane over the course of these years; Reinhard, the fastest postdoc at DØ , and Bob H., who showed me the artsy side of Chicago; Juan and Kyle, for being such good friends; Paul and Randal, for stimulating discussions about particle physics and beyond; Aneta and Richard, for helping me find my way through the jungle of UIC thesis regulations; and Jui, for her perfect assistance as UIC graduate adviser.

I could not have survived without support from the UIC “gang”. Thanks Gustavo, James,

ACKNOWLEDGMENTS (Continued)

and Tim, for some unforgettable moments. Special thanks to Mario for introducing me to the insanely fine art of Espresso.

Mom and dad, thank you for your patience and support, while I was pursuing my studies far away from home. Thanks to Anja, for being a great sister.

Finally, I would like to thank Dasha for her love. This dissertation would not have been possible without her encouragement.

MB

TABLE OF CONTENTS

<u>CHAPTER</u>	<u>PAGE</u>
1 INTRODUCTION	1
2 THEORY	3
2.1 The Standard Model	3
2.1.1 Quarks and Leptons	3
2.1.2 Interactions	6
2.1.2.1 Electromagnetic Interaction	6
2.1.2.2 Weak Interaction	7
2.1.2.3 Strong Interaction	9
2.2 Cross Sections	11
2.3 The Measurement of the $Z/\gamma^*(\rightarrow e^+e^-) + \geq n$ Jet Cross Sections	14
3 EXPERIMENTAL APPARATUS	18
3.1 The Fermilab Accelerators	18
3.1.1 The Preaccelerator	20
3.1.2 The Linac	21
3.1.3 The Booster	22
3.1.4 The Main Injector	23
3.1.5 The Antiproton Source	23
3.1.6 The Tevatron	24
3.2 Luminosity and Cross Section	25
3.3 The DØ Detector	27
3.3.1 Coordinate Systems	27
3.3.2 Central Tracking System	30
3.3.2.1 Silicon Microstrip Tracker	30
3.3.2.2 Central Fiber Tracker	32
3.3.2.3 Solenoidal Magnet	35
3.3.3 Preshower Detectors	35
3.3.3.1 Central Preshower Detector	36
3.3.3.2 Forward Preshower Detector	38
3.3.4 Calorimeter	39
3.3.5 Muon System	41
4 THE DØ TRIGGER AND DATA ACQUISITION SYSTEMS .	46
4.1 The Level 1 Trigger	46
4.1.1 The Level 1 Calorimeter Trigger	47

TABLE OF CONTENTS (Continued)

<u>CHAPTER</u>		<u>PAGE</u>
	4.1.2 The Level 1 Central Track Trigger	47
	4.1.3 The Level 1 Muon Trigger	47
	4.2 The Level 2 Trigger	48
	4.2.1 The Level 2 Calorimeter Preprocessor	49
	4.2.2 The Level 2 Muon Preprocessors	50
	4.2.3 The Level 2 Preshower Preprocessor	50
	4.2.4 The Level 2 Tracking Preprocessor	50
	4.2.5 The Level 2 Global Processor	51
	4.3 The Level 3 Trigger and Data Acquisition	52
5	OFFLINE EVENT RECONSTRUCTION	53
	5.1 Track Reconstruction	53
	5.2 Primary Vertex Reconstruction	54
	5.3 Electromagnetic Object Reconstruction and Identification . .	56
	5.3.1 Simple-Cone Clustering Algorithm	57
	5.3.2 Electromagnetic Object Identification Parameters	57
	5.4 Jet Reconstruction and Identification	60
	5.4.1 Jet Cone Algorithm	60
	5.4.2 Jet Identification Parameters	62
	5.4.3 Jet Energy Scale	64
6	MONTE CARLO EVENT SIMULATION	66
	6.1 The PYTHIA Event Generator	66
	6.2 Combining Matrix Elements with Showering	68
	6.2.1 Combining ALPGEN with PYTHIA	70
	6.2.2 Combining MADGRAPH with PYTHIA	70
	6.3 The MCFM Monte Carlo Simulation	71
	6.4 The DØ Detector Simulation	71
7	DATA AND MONTE CARLO SAMPLES	72
	7.1 Data Sample	72
	7.2 Monte Carlo Samples	74
	7.2.1 PYTHIA and ALPGEN Samples	74
	7.2.2 CKKW Samples	77
	7.2.3 MCFM Cross Sections	79
	7.3 Event Selection	79
	7.3.1 Primary Vertex	79
	7.3.2 Electron Selection	79
	7.3.3 Z Selection	80
	7.3.4 Jet Selection	80
	7.3.5 Event Statistics	81
	7.4 Data vs Monte Carlo	81

TABLE OF CONTENTS (Continued)

<u>CHAPTER</u>		<u>PAGE</u>
7.4.1	Primary Vertex Comparison	82
7.4.2	Z p_T Comparisons	82
7.4.3	Z/γ^* ($\rightarrow e^+e^-$) + $\geq n$ Jet Comparisons	87
7.4.3.1	Z/γ^* ($\rightarrow e^+e^-$) Inclusive Sample	87
7.4.3.2	Z/γ^* ($\rightarrow e^+e^-$) + ≥ 1 Jet Sample	87
7.4.3.3	Z/γ^* ($\rightarrow e^+e^-$) + ≥ 2 Jet Sample	87
8	MEASUREMENT OF THE $Z/\gamma^*(\rightarrow e^+e^-)$ INCLUSIVE CROSS SECTION	99
8.1	Efficiencies	100
8.1.1	Trigger Efficiency	100
8.1.2	EM Reconstruction and Identification Efficiency	102
8.1.3	EM-Track Match Efficiency	105
8.1.4	Acceptance	108
8.2	Cross Section Calculation	111
8.3	Comparisons to Other Measurements	113
9	MEASUREMENT OF THE $Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ JET CROSS SECTIONS	116
9.1	Efficiencies vs Jet Multiplicity	116
9.1.1	Trigger Efficiency	116
9.1.2	EM Reconstruction and Identification Efficiency	117
9.1.3	EM-Track Match Efficiency	118
9.1.4	Acceptance	122
9.1.5	Jet Reconstruction and Identification Efficiency	123
9.2	Cross Section Calculation	126
9.2.1	Unsmearing	126
9.2.2	Electron-Jet-Overlap Correction	138
9.2.3	Cross Sections	144
10	SYSTEMATICS	151
10.1	Jet Energy Scale Systematic Error	151
10.2	Systematic Error of Cross Section Unfolding	151
10.3	Electron-Jet-Overlap Systematic Error	154
10.4	Luminosity Systematic Error	161
10.5	Systematic Errors Due to Efficiencies	162
10.5.1	Trigger Efficiency	162
10.5.2	EM Reconstruction and Identification Efficiency	163
10.5.3	EM-Track Match Efficiency	163
10.5.4	Overall Efficiency Systematic Error	164
10.6	Jet Promotion Systematic Error	164
10.7	Statistical Uncertainty	165

TABLE OF CONTENTS (Continued)

<u>CHAPTER</u>		<u>PAGE</u>
11	CONCLUSIONS	170
11.1	Results	170
11.2	Future Plans	171
	APPENDICES	176
	Appendix A	177
	Appendix B	191
	CITED LITERATURE	200
	VITA	207

LIST OF TABLES

<u>TABLE</u>		<u>PAGE</u>
I	SINGLE EM TRIGGERS USED IN THIS ANALYSIS	75
II	LIST OF MONTE CARLO SAMPLES	77
III	JET ENERGY RESOLUTION PARAMETERS IN DATA AND MC.	78
IV	LIST OF CKKW SAMPLES	78
V	EVENT BREAKDOWN BY EXCLUSIVE JET MULTIPLICITIES ASSOCIATED WITH Z/γ^* PRODUCTION BEFORE ANY BACK- GROUND IS SUBTRACTED OR ANY CORRECTIONS ARE AP- PLIED.	82
VI	CROSS SECTION RESULTS FOR DIFFERENT DIEM INVARI- ANT MASS RANGES AND PROCESSES.	115
VII	OBJECT BASED TRIGGER EFFICIENCIES WITH STATISTI- CAL AND SYSTEMATIC ERRORS (RESPECTIVELY) FOR THE PRE-V12 AND V12 DATASETS FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES.	118
VIII	OBJECT BASED EM RECO AND ID EFFICIENCIES WITH STA- TISTICAL AND SYSTEMATIC ERRORS (RESPECTIVELY) IN DATA AND MC FOR DIFFERENT INCLUSIVE JET MULTIPLI- CITIES. THERE WAS NOT ENOUGH STATISTICS AVAILABLE TO ESTIMATE THE EM EFFICIENCY IN DATA FOR ≥ 3 JETS.	120
IX	OBJECT BASED TRACKING EFFICIENCIES WITH STATISTI- CAL ERRORS IN DATA AND MC FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES.	120
X	OBJECT BASED TRACKING EFFICIENCIES WITH SYSTEM- ATIC ERRORS.	122
XI	ACCEPTANCES WITH STATISTICAL ERRORS FOR DIFFER- ENT JET MULTIPLICITIES.	123

LIST OF TABLES (Continued)

<u>TABLE</u>		<u>PAGE</u>
XII	UNSMEARING AND JET RECO/ID COEFFICIENTS WITH SYSTEMATIC UNCERTAINTY DUE TO RESOLUTION AND JET RECO/ID EFFICIENCY.	137
XIII	ELECTRON-JET-OVERLAP COEFFICIENTS WITH SYSTEMATIC UNCERTAINTIES.	140
XIV	NUMBER OF SIGNAL EVENTS (DRELL-YAN IN PARENTHESES) AND NUMBER OF BACKGROUND EVENTS (QCD) FOR DIFFERENT JET MULTIPLICITIES. NUMBERS ARE CORRECTED FOR TRIGGER, EM AND TRACKING INEFFICIENCIES (AFTER APPLYING THE UNSMEARING, JET RECO/ID AND ELECTRON-JET-OVERLAP CORRECTIONS).	144
XV	FINAL CROSS SECTIONS WITH JET ENERGY SCALE ERRORS.	153
XVI	FINAL CROSS SECTIONS WITH JET RECO/ID ERRORS. . . .	154
XVII	FINAL CROSS SECTIONS WITH JET RESOLUTION ERRORS.	161
XVIII	FINAL CROSS SECTIONS WITH ELECTRON-JET-OVERLAP CUT UNCERTAINTIES.	161
XIX	FINAL CROSS SECTIONS WITH LUMINOSITY ERRORS.	162
XX	RELATIVE ERRORS DUE TO UNCERTAINTY IN EM-TRACK MATCHING EFFICIENCIES.	164
XXI	OVERALL SYSTEMATIC UNCERTAINTIES DUE TO EFFICIENCIES (TRIGGER, EM, TRACKING).	165
XXII	NUMBER OF EVENTS FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES WHEN REQUIRING EXACTLY ONE RECONSTRUCTED PRIMARY VERTEX AND AT LEAST TWO RECONSTRUCTED PRIMARY VERTICES. ENTRIES ARE NORMALIZED WITH RESPECT TO THE 2 VERTEX SAMPLE.	166
XXIII	AVERAGE NUMBER OF RECONSTRUCTED PRIMARY VERTICES FOR DIFFERENT JET MULTIPLICITIES.	166
XXIV	CROSS SECTIONS WITH STATISTICAL ERRORS.	169

LIST OF TABLES (Continued)

<u>TABLE</u>		<u>PAGE</u>
XXV	CROSS SECTIONS FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES. NUMBER OF SIGNAL EVENT ENTRIES HAVE UNSMEARING, JET RECO/ID AND ELECTRON-JET-OVERLAP CORRECTIONS APPLIED.	171
XXVI	COMPARISON OF MEASURED CROSS SECTIONS WITH RESULTS FROM MCFM AND CKKW.	172
XXVII	COMPARISON OF MEASURED CROSS SECTION RATIOS WITH RESULTS FROM MCFM AND CKKW.	172

LIST OF FIGURES

<u>FIGURE</u>	<u>PAGE</u>
1 Properties of fermion particles: quarks and leptons. Spin is given in units of \hbar , electric charge is given in units of the absolute value of the electron charge.	5
2 Fundamental forces and their properties.	7
3 Feynman diagram for LO Z production via $q\bar{q} \rightarrow Z \rightarrow e^+e^-$	13
4 Feynman diagram for Z/γ^* production in association with a parton via $q\bar{q} \rightarrow Z/\gamma^*g$	15
5 Feynman diagrams for Z/γ^* production in association with a parton via $qg \rightarrow Z/\gamma^*q$	16
6 SM Higgs boson production in association with a Z boson via $q\bar{q} \rightarrow HZ$ (<i>Higgsstrahlung</i>).	16
7 Feynman diagram for Z/γ^* production in association with two gluons via $q\bar{q} \rightarrow Z/\gamma^*gg$	17
8 Schematic view of the Fermilab accelerator chain.	19
9 Schematic view of magnetron operation for the hydrogen ion source. . .	21
10 Schematic of Linac RF cavity.	22
11 Simplified drawing of anti-proton production with nickel target and lithium lens.	24
12 Tevatron integrated luminosity delivered to DØ (April 2002 - June 2005). The arrow indicates the data taking period during which the data for this analysis was recorded (343 pb^{-1}).	26
13 Schematic view of the DØ detector.	28
14 The DØ central tracking system with with solenoid, preshower detectors, luminosity monitor, and calorimeter.	31

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
15	Double-sided ladder design, n-side.	32
16	SMT disk and barrel design.	33
17	a) Location of the Central Fiber Tracker (CFT). b) Closeup view of axial and stereo layers.	34
18	$y - z$ view of the DØ magnetic field with both the toroid and solenoid magnets at full current. Numbers are in kG (10 kG = 1 T).	36
19	Perspective view of the solenoid inside the central calorimeter.	37
20	Cross section and layout geometry of the CPS and FPS scintillator strips.	38
21	Isometric view of the central and two end calorimeters.	41
22	Schematic view of two calorimeter cells.	42
23	Schematic view showing the calorimeter segmentation pattern. The shading pattern indicates cells for signal readout. The radial lines show the detector pseudo-rapidity intervals.	43
24	Schematic view of different calorimeter detection layers vs η	44
25	The DØ muon system.	45
26	A hypothetical L1CTT track with hits in eight CFT axial doublet layers and the CPS axial layer.	48
27	The definition of roads based on L1 tracks and SMT hit selection in L2STT.	51
28	Axial view (looking down the beam-pipe) of a recorded event showing hits and reconstructed tracks. CFT hits are represented by squares, and SMT hits are represented by circles. Hits are colored solid if they are associated with a reconstructed track (solid lines). The curvature of the reconstructed tracks is due to the solenoidal magnetic field, which is pointing out of the page.	55
29	Illustration showing how the presence of soft radiation between two jets may cause a merging of the jets (right) that would not occur in the absence of the soft radiation (left).	62

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
30	The jet energy scale correction factor measured for jets in data as a function of E (top) and η (bottom).	65
31	Schematic view of a parton shower.	69
32	Primary vertex distribution in data and MC (PYTHIA) for the inclusive sample. The MC distribution is normalized to the number of events in data.	83
33	Comparison of Z p_T between data and PYTHIA MC (left), and ratio correction factor (right) for the inclusive sample. The MC distribution (left) is normalized to the number of events in data.	84
34	Comparison of Z p_T between data and ALPGEN + PYTHIA $Z+1$ jet MC. The MC distribution is normalized to the number of events in data.	85
35	Comparison of Z p_T between data and ALPGEN + PYTHIA $Z+2$ jets MC. The MC distribution is normalized to the number of events in data.	86
36	p_T of both Z electrons (top left), physics η of both Z electrons (bottom left), Z p_T (top right), Z rapidity (bottom right) for the $Z/\gamma^* \rightarrow e^+e^-$ inclusive sample in data and MC (PYTHIA). The MC distribution is normalized to the number of events in data.	88
37	Diem invariant mass comparison for the $Z/\gamma^* \rightarrow e^+e^-$ inclusive sample in data and MC (PYTHIA). Data is background subtracted. The MC distribution is normalized to the number of events in data.	89
38	p_T of both Z electrons (top left), physics η of both Z electrons (bottom left), Z p_T (top right), Z rapidity (bottom right) for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.	90
39	Diem invariant mass comparison for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample in data and MC (ALPGEN). Data is background subtracted. The MC distribution is normalized to the number of events in data.	91
40	p_T (linear and logarithmic), physics η and physics Φ of all jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.	92

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
41	p_T (linear and logarithmic), physics η and physics Φ of the leading jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data. . . .	93
42	p_T of both Z electrons (top left), physics η of both Z electrons (bottom left), Z p_T (top right), Z rapidity (bottom right) for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.	94
43	Diem invariant mass comparison for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). Data is background subtracted. The MC distribution is normalized to the number of events in data.	95
44	p_T (linear and logarithmic), physics η and physics Φ of all jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.	96
45	p_T (linear and logarithmic), physics η and physics Φ of the leading jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data. . . .	97
46	p_T (linear and logarithmic), physics η and physics Φ of the next to leading jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.	98
47	Trigger efficiencies for pre-v12 (top) and v12 (bottom) datasets vs EM object p_T	102
48	EM efficiencies versus probe track Φ and p_T in data. The Φ distribution shows the modulus($\Phi, \frac{2\pi}{32}$) distribution to illustrate the effect of the calorimeter Φ -module boundaries.	106
49	EM efficiencies versus probe track Φ and p_T in MC. The Φ distribution shows the modulus($\Phi, \frac{2\pi}{32}$) distribution to illustrate the effect of the calorimeter Φ -module boundaries.	107
50	Invariant mass in data when requiring at least one track-matched electron.	109
51	Invariant mass in data when requiring two track-matched electrons. . .	109
52	Invariant mass in MC when requiring at least one track-matched electron.	110

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
53	Invariant mass in MC when requiring two track-matched electrons. . . .	110
54	Diem invariant mass distribution for $Z/\gamma^* \rightarrow e^+e^- + X$ (Mean = 91.02 GeV \pm 0.04 GeV, Width 4.03 GeV \pm 0.04 GeV).	114
55	Average object based trigger efficiencies in data versus inclusive jet multiplicity.	117
56	Average object based EM reco and ID efficiencies in data and MC versus inclusive jet multiplicity. There was not enough statistics available to estimate the EM efficiency in data for ≥ 3 jets.	119
57	Average object based tracking efficiencies in data and MC versus inclusive jet multiplicity.	121
58	Jet reco/ID efficiencies in data. CC = $-0.7 < \eta_{det} < 0.7$, ICR = $0.7 < \eta_{det} < 1.5$, EC = $1.5 < \eta_{det} < 2.5$	125
59	Comparing inclusive jet multiplicities between data and particle level MC (applying data resolution smearing and data jet reco/ID efficiencies). The distributions are normalized with respect to the first bin.	127
60	Ratio of MC (with smearing and jet reco/ID efficiencies) inclusive jet multiplicities and data inclusive jet multiplicities.	128
61	Comparing inclusive jet multiplicities between data and particle level MC (after applying correction factors). The distributions are normalized with respect to the first bin.	128
62	Comparing jet p_T for all jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied).	129
63	Comparing jet p_T for all jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied).	129
64	Comparing jet p_T for leading jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.	130

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
65	Comparing jet p_T for leading jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.	130
66	Comparing jet p_T for second leading jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.	131
67	Comparing jet p_T for second leading jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.	131
68	Comparing jet p_T for third leading jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied).	132
69	Comparing jet p_T for third leading jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied).	132
70	Comparing jet η for all jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied).	133
71	Comparing jet η for leading jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied). . .	133
72	Comparing jet η for second leading jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied). . .	134
73	Comparing jet η for third leading jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied). . .	134
74	Unsmearing and jet reco/ID particle jet multiplicities (left) and coefficients (right).	135
75	Unsmearing jet multiplicities (left) and coefficients (right) without applying jet reco/ID efficiencies.	136

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
76	ΔR between probe tracks and good jets in data (without electron-jet-overlap cut).	139
77	ΔR between probe tracks and good jets using PYTHIA MC (without electron-jet-overlap cut).	140
78	ΔR between probe tracks and good jets in data (after the electron-jet-overlap cut was applied).	141
79	ΔR between generated electrons ($p_T > 25$ GeV, $ \eta < 1.1$) and partons ($p_T > 20$ GeV, $ \eta < 2.5$).	142
80	ΔR between partons and matched calorimeter jets ($p_T > 20$ GeV, $ \eta < 2.5$).	143
81	Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample. The solid line shows a Gaussian plus Breit-Wigner fit to the Z peak. The dashed line shows an exponential fit to the QCD and Drell-Yan contribution.	145
82	Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample. The solid line shows a Gaussian plus Breit-Wigner fit to the Z peak. The dashed line shows an exponential fit to the QCD and Drell-Yan contribution.	146
83	Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 3$ jet sample.	147
84	Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 4$ jet sample.	148
85	Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 5$ jet sample.	149
86	$Z/\gamma^*(\rightarrow e^+e^-) + \geq n$ jet cross sections with statistical errors.	150
87	$\pm 1\sigma$ fluctuation of the jet energy scale (JES 5.3)	152
88	Jet reco/ID efficiencies with errors plotted versus particle jet p_T smeared with data energy resolution (central).	155
89	Jet reco/ID efficiencies with errors plotted versus particle jet p_T smeared with data energy resolution (ICR).	156
90	Jet reco/ID efficiencies with errors plotted versus particle jet p_T smeared with data energy resolution (forward).	157

LIST OF FIGURES (Continued)

<u>FIGURE</u>		<u>PAGE</u>
91	Comparing scale factors between the Top group and this analysis. Equivalent techniques were used in both cases.	158
92	Comparing “efficiency” of finding a recoiling jet as a function of Z p_T in data and MC.	159
93	Comparing jet resolution for JES 5.1 (upper curve) and JES 5.3 (lower curve) in the central region of the Calorimeter. The difference is approximately 5% over the whole range.	160
94	Exponential fit to the number of QCD events for different inclusive jet multiplicities.	168
95	$Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ jet cross sections in data (with total errors) compared with MCFM and CKKW.	173
96	Ratios of the $Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ jet cross sections to the total inclusive $Z/\gamma^* \rightarrow e^+e^-$ cross section versus n . The errors on the data include the combined statistical and systematic uncertainties. The dashed line (CKKW) represents the predictions of LO matrix element calculations using PYTHIA for parton showering and hadronization, normalized to the measured $Z/\gamma^*+\geq 1$ jet cross section ratio. The diamonds represent the MCFM predictions.	174
97	Data to theory (ALPGEN+PYTHIA) comparison for the highest p_T jet distribution in the $Z+\geq 1$ jet sample, for the second highest p_T jet distribution in the $Z+\geq 2$ jet sample and for the third highest p_T jet distribution in the $Z+\geq 3$ jet sample. The errors on the data are only statistical.	175
98	Schematic view of subdetectors with L1 and L2 trigger elements. Horizontal arrows indicate the direction of dataflow.	192
99	Layout of Level 2 calorimeter (L2CAL) crate with main components. .	195
100	Level 2 Alpha Board with Ethernet card and hard disk drive.	196
101	Block diagram of Level 2 Alpha Board.	197
102	Block diagram of the 21172 Core Logic Chipset.	199

LIST OF ABBREVIATIONS

AA	Alternative Algorithm
BR	Branching Ratio
CC	Central Calorimeter
CellNN	Cell Nearest Neighbor
CDF	Collider Detector at Fermilab
CERN	European Organization for Nuclear Research
CFT	Central Fiber Tracker
CH(F)	Coarse Hadronic (Fraction)
CIA	Control, Input/Output, Address
CKKW	Catani, Krauss, Kuhn, Webber
COOR	Central Coordination Process
CPLD	Complex Programmable Logic Device
CPS	Central Preshower
CPU	Central Processing Unit
CTEQ	Coordinated Theoretical-Experimental Project on QCD

LIST OF ABBREVIATIONS (Continued)

D0GSTAR	D0 GEANT Simulation of the Total Apparatus Response
DAQ	Data Acquisition
DEC	Digital Equipment Corporation
DGLAP	Dokshitzer-Gribov-Lipatov-Altarelli-Parisi
DIS	Deep Inelastic Scattering
DMA	Direct Memory Access
DRAM	Dynamic Random Access Memory
DSW	Data Switch
EC	End Calorimeter
EM	Electro Magnetic
EMF	Electro Magnetic Fraction
FAMUS	Forward Angle Muon System
FIC	Fiber Input Converter
FH	Fine Hadronic
FIFO	First in, First out
FNAL	Fermi National Accelerator Laboratory
FPGA	Field Programmable Gate Array
FPS	Forward Preshower

LIST OF ABBREVIATIONS (Continued)

FSR	Final State Radiation
GEANT	GEometry ANd Tracking
GWS	Glashow, Weinberg, Salam
HotF	Hot Fraction
HTF	Histogramming Track Finder
ICD	Inter Cryostat Detector
ICR	Inter Cryostat Region
ISR	Initial State Radiation
IDE	Integrated Drive Electronics
I/O	Input/Output
ISA	Industry Standard Architecture
JES	Jet Energy Scale
L1	Level 1
L1Cal	Level 1 Calorimeter
L1CTT	Level 1 Central Track Trigger
L1Muon	Level 1 Muon
L2	Level 2
L2Cal	Level 2 Calorimeter

LIST OF ABBREVIATIONS (Continued)

L2CTT	Level 2 Central Track Trigger
L2GBL	Level 2 Global
L2MUC	Level 2 Muon Central
L2MUF	Level 2 Muon Forward
L2Muon	Level 2 Muon
L2PS	Level 2 Preshower
L2STT	Level 2 Silicon Track Trigger
L3	Level 3
L3DAQ	Level 3 Data Acquisition
LBN	Luminosity Block Number
Linac	Linear Accelerator
LL	Leading Logarithmic
LLA	Leading Log Approximation
MB	Minimum Bias
MBT	Magic Bus Transceiver
MBus	Magic Bus
MC	Monte Carlo
MCFM	Monte Carlo for FeMtobarn processes

LIST OF ABBREVIATIONS (Continued)

MDT	Mini Drift Tube
MG	Massless Gap
MPM	Multi Port Memory
NLO	Next-To-Leading Order
PCI	Peripheral Component Interconnect
PDF	Parton Distribution Function
PDT	Proportional Drift Tube
PIO	Programmed Input Output
PLD	Programmable Logic Device
PV	Primary Vertex
RF	Radio Frequency
RISC	Reduced Instruction Set Computing
ROM	Read Only Memory
QCD	Quantum Chromo Dynamics
QED	Quantum Electro Dynamics
QFT	Quantum Field Theory
SBC	Single Board Computer
SCL(init)	Serial Command Link (Initialize)

LIST OF ABBREVIATIONS (Continued)

Scone	Simple Cone
SIMM	Single In-line Memory Module
SLIC	Second Level Input Computer
SM	Standard Model
SMT	Silicon Microstrip Tracker
SRAM	Serial Read Only Memory
TCC	Trigger Control Computer
TDR	Technical Design Report
TOT	Total Energy
VBD	VME Buffer Driver
VITA	VMEbus International Trade Association
VLPC	Visible Light Photon Counter
VME	VERSA Module Eurocard
WAMUS	Wide Angle Muon System

SUMMARY

A study of events with Z/γ^* bosons and hadronic jets produced at the Tevatron in $p\bar{p}$ collisions at a center of mass energy of 1.96 TeV is presented. The data consist of approximately 14,000 $Z/\gamma^* \rightarrow e^+e^-$ decay candidates from 343 pb^{-1} of integrated luminosity collected using the DØ detector. Cross sections and jet production properties have been measured for $Z/\gamma^* + \geq 0$ to 5 jet events. This measurement represents a significant improvement over previous measurements at the Tevatron. It is the first measurement at this center of mass energy with the DØ detector. The results are in good agreement with QCD predictions.

CHAPTER 1

INTRODUCTION

Since the discovery of the Z boson by the European Organization for Nuclear Research (CERN) in 1983 (1; 2), the production of electroweak gauge bosons in association with jets has been of increasing importance. A measurement of the $Z + \geq n$ jet cross sections provides a good test of the current understanding of quantum chromo dynamics (QCD). In addition, the associated production of Z bosons with jets is a major background to many interesting physics processes. In particular, searches in the channel where a Higgs boson is being produced in association with a Z boson rely on a detailed understanding of the $Z + \geq n$ jet background.

Chapter 2 of this dissertation provides a short overview of the theoretical framework within which this analysis was performed. A description of the experimental apparatus, including the Fermilab chain of accelerators, and the DØ detector is given in Chapter 3. Chapter 4 describes the methods that are used to reconstruct physical objects from raw detector data. The Monte Carlo (MC) event generators that are used are illustrated in Chapter 5. Data and MC samples, including comparisons between data and MC, are described in Chapter 6. Chapters 7 and 8 outline all steps that lead to the measurements of the inclusive $Z/\gamma^*(\rightarrow e^+e^-)$ and $Z/\gamma^*(\rightarrow e^+e^-) + \geq n$ jet cross sections. Sources for systematic uncertainties are discussed in Chapter 9. A summary of the results, together with a discussion of future plans, is given in Chapter 10.

Throughout this dissertation (unless stated otherwise) the customary convention by which $\hbar = c = 1$ is followed.

CHAPTER 2

THEORY

2.1 The Standard Model

The *Standard Model (SM) of Particle Physics* is the current theory of elementary particles and the forces that act between them. The SM is a quantum theory of fields (QFT), which arises when combining quantum mechanics with special relativity. The SM includes most of the current understanding of the laws of physics (except gravity), and has been verified experimentally to an extremely high level of accuracy.

Nevertheless, the theory is incomplete. The SM contains many free parameters that cannot be derived from first principles. The Higgs boson, which is considered to be the last remaining piece to the SM, has not been experimentally detected yet. Most importantly, gravity is not included in the SM. A framework that consistently combines quantum mechanics with general relativity into a theory of *quantum gravity* has not been experimentally confirmed yet.

The following gives an overview of the SM (3; 4; 5; 6; 7; 8; 9).

2.1.1 Quarks and Leptons

In the SM the fundamental particles that make up ordinary matter are divided into two groups: *quarks* and *leptons* (Figure 1, Reference (10)). Both quarks and leptons are *fermions* since they are spin- $\frac{1}{2}$ particles, and therefore obey Fermi-Dirac statistics. As indicated in Figure 1, quarks and leptons are each arranged in three generations, containing particles of

similar properties but differing mass. For each particle there exists an associated anti-particle, which is the same as its corresponding particle in properties such as mass and spin, and opposite in properties such as charge and magnetic moment.

There are six different flavors of quarks, labeled (in order of increasing mass) *up*, *down*, *strange*, *charm*, *bottom*, and *top*. It is remarkable to note that quarks are never observed as single particles (see Chapter 2.1.2.3), and that they carry fractional electrical charges¹ of $+\frac{2}{3}$ or $-\frac{1}{3}$. Quarks form bound states called *hadrons* by either combining three quarks into *baryons*, or by pairing a quark with an antiquark into *mesons*. Protons (made up of two up-quarks and one down-quark) and neutrons (made up of two down-quarks and one up-quark) are the most common examples of baryons. Pions ($\pi^{0,\pm}$) and Kaons (K^\pm , K^0 , \bar{K}^0) are the most common type for mesons.

There are three different flavors of charged leptons carrying a charge of -1: *electron* (e^-), *muon* (μ^-), and *tau* (τ^-). While electrons exist in all atoms, muons and taus can only be observed in energetic processes like cosmic ray showers, or high energy particle collisions. There are three neutral leptons, called *neutrinos* (ν), each corresponding to a charged lepton: ν_e , ν_μ , and ν_τ . Neutrinos interact extraordinarily weakly with matter, and their masses are negligibly small.

¹All charges are given in units of the absolute value of the electron charge, 1.602×10^{-19} Coulombs.

Leptons spin = 1/2			Quarks spin = 1/2		
Flavor	Mass GeV/c ²	Electric charge	Flavor	Approx. Mass GeV/c ²	Electric charge
ν_e electron neutrino	$<1 \times 10^{-8}$	0	u up	0.003	2/3
e electron	0.000511	-1	d down	0.006	-1/3
ν_μ muon neutrino	<0.0002	0	C charm	1.3	2/3
μ muon	0.106	-1	S strange	0.1	-1/3
ν_τ tau neutrino	<0.02	0	t top	175	2/3
τ tau	1.7771	-1	b bottom	4.3	-1/3

Figure 1. Properties of fermion particles: quarks and leptons. Spin is given in units of \hbar , electric charge is given in units of the absolute value of the electron charge.

2.1.2 Interactions

One of the most fundamental insights in theoretical physics is that interactions are dictated by symmetry principles. In QFT, interactions manifest themselves by imposing symmetry conditions on the quantum fields representing the respective interactions. Using the framework of Lagrangian field theory, the Lagrangian of the theory is required to be invariant under a group of local phase changes (*local gauge invariance*). A local phase depends on space and time in a completely arbitrary way. The Lagrangian is no longer invariant under such local phase changes. In order to preserve gauge invariance of the Lagrangian (on purely aesthetic grounds), an additional gauge field is introduced. Excitations of these gauge fields lead to spin-1 bosons that mediate the interactions of nature.

Four types of interactions are currently known (in order of decreasing strength): strong, electromagnetic, weak, and gravitational. Figure 2 summarizes basic properties of the four interactions (10). Note that gravity has not been incorporated into the SM, since no satisfactory QFT for gravitational interactions has been established yet. Additionally, gravity is simply too weak to have any significant influence on particle interactions.

The following is a brief summary of the interactions that have been incorporated into the SM.

2.1.2.1 Electromagnetic Interaction

Historically, the electromagnetic interaction was the first to be formulated in the framework of a renormalizable QFT by Tomonaga, Feynman, and Schwinger in the 1940s (Nobel Prize in 1965). *Quantum Electrodynamics* (QED) describes the electromagnetic interaction by

Property \ Interaction	Gravitational	Weak (Electroweak)	Electromagnetic	Strong	
				Fundamental	Residual
Acts on:	Mass – Energy	Flavor	Electric Charge	Color Charge	See Residual Strong Interaction Note
Particles experiencing:	All	Quarks, Leptons	Electrically charged	Quarks, Gluons	Hadrons
Particles mediating:	Graviton (not yet observed)	$W^+ W^- Z^0$	γ	Gluons	Mesons
Strength relative to electromag for two u quarks at: for two protons in nucleus	10^{-41}	0.8	1	25	Not applicable to quarks
	10^{-41}	10^{-4}	1	60	
	10^{-36}	10^{-7}	1	Not applicable to hadrons	20

Figure 2. Fundamental forces and their properties.

requiring gauge invariance under $U(1)$ group transformations. $U(1)$ denotes a group of unitary one-dimensional matrices, describing space-time dependent rotations in a complex plane. The requirement of gauge invariance gives rise to the photon field and the *photons* as the corresponding mediator of the electromagnetic interaction. Because the photon is massless the interaction is long range. The photon couples to all particles that carry electrical charge, like quarks and charged leptons. The strength of the interaction is proportional to the magnitude of the dimensionless *fine structure coupling constant*:

$$\alpha_{EM} = \frac{e^2}{4\pi} \approx \frac{1}{137} \quad (2.1)$$

2.1.2.2 Weak Interaction

The weak interaction is most prominent in *beta decays* (Equation 2.2 and Equation 2.3) and associated radioactivity.

$$n \rightarrow p + e^- + \bar{\nu}_e \quad (2.2)$$

$$p \rightarrow n + e^+ + \nu_e \quad (2.3)$$

The range of the interaction is short due to the high mass of the mediating gauge bosons (W^\pm , Z^0):

$$m_{W^\pm} = 80.425 \pm 0.038 \text{ GeV} \quad (2.4)$$

$$m_{Z^0} = 91.1876 \pm 0.0021 \text{ GeV} \quad (2.5)$$

A QFT combining the electromagnetic with the weak interaction was first developed by Glashow, Weinberg, and Salam (*GWS theory*, Noble Prize in 1979). Later 't Hooft and Veltman were able to prove that the theory is renormalizable (Nobel Prize in 1999). Electroweak theory combines a new U(1) group with an SU(2) group, and requires invariance under $SU(2) \otimes U(1)$ transformations. SU(n) describes groups of special¹ and unitary $n \times n$ matrices. Local gauge invariance under SU(2) group transformations introduces three massless spin-1 gauge bosons W^+ , W^- , and W^0 . Adding the U(1) group introduces another gauge boson called B^0 . The W^0 and B^0 mix quantum mechanically to give rise to the experimentally observed photon (γ) and Z^0 :

$$\gamma = W^0 \sin \theta_W + B^0 \cos \theta_W \quad (2.6)$$

$$Z^0 = W^0 \cos \theta_W - B^0 \sin \theta_W \quad (2.7)$$

¹The determinant of the matrices must be 1.

where θ_W is called the *weak mixing angle* or *Weinberg angle*. As opposed to QED, the underlying group of the electroweak theory is *non-Abelian* since not all the generators of the group commute with each other. This ultimately leads to self interaction between the gauge bosons.

Up to this point the electroweak theory is very simple and elegant. Yet it is incomplete, since all particles of the theory are massless. However, it is known from experiment that the W and Z bosons are heavy. Additionally, mass terms cannot be introduced into the Lagrangian describing the system, since this would destroy the gauge invariance of the Lagrangian. This problem is resolved by the *Higgs mechanism*, which introduces *spontaneous symmetry breaking* of the Higgs scalar field potential, thereby giving mass to the gauge bosons (W and Z) and the quark and leptons.

2.1.2.3 Strong Interaction

Quantum Chromo Dynamics (QCD) is the QFT describing the strong interaction. It is based on an $SU(3)$ gauge field, which leads to 8 mediating massless gauge bosons called *gluons*. Strongly interacting particles like quarks and gluons carry a new type of charge called *color*. Each (anti)quark can carry a (anti)red, (anti)green, or (anti)blue color charge. Gluons carry a combination of a color and anticolor charge. The fact that gluons are themselves carriers of the color charge allows them to couple to each other. This derives from the non-Abelian character of the gauge theory. Quarks and gluons are collectively referred to as *partons*.

One interesting feature of QCD is the fact that the strength of the strong coupling increases with decreasing energy scale, i.e. at low energies and long distances the interaction becomes

too strong to be treated within the framework of perturbation theory. This leads to the idea of *confinement*, which assumes that all objects carrying color can never be found as free particles in nature and that they are confined into color-neutral composite hadrons. Colors can be combined into color-neutral objects in two ways: a colored quark combined with an antiquark of the corresponding anticolor (meson), or a combination of three (anti)quarks each carrying a different (anti)color (baryon). This confines quarks to a small region near one another ($\approx 10^{-15}$ m). The quarks that combine into baryons or mesons are referred to as *valence quarks*. They constantly interact with each other by exchanging gluons. Since gluons can couple to each other, they can emit more gluons that can further split into virtual quark-antiquark pairs called *sea quarks*.

Experimentally quarks and gluons are observed as *jets* of color-neutral hadrons. This means that if a single parton emerges from a particle collision, gluons will be radiated which subsequently produce quark-antiquark pairs to form a *parton shower*. Ultimately the partons combine into a jet of hadrons moving in the direction close to that of the original parton. This final step is called *hadronization*.

Mathematically the strong coupling constant, α_s , takes the approximate form:

$$\alpha_s(Q^2) \approx \frac{12\pi}{(11c - 2n_f) \log(\frac{Q^2}{\Lambda^2})}, \quad (2.8)$$

where Q expresses the magnitude of the momentum transferred in the interaction, n_f indicates the number of quarks (6 in the SM), and c is the number of quark colors (3 in the SM). Λ is the QCD scale parameter, defined as:

$$\Lambda^2 = \mu_R^2 \exp \frac{-12\pi}{(11c - 2n_f)\alpha_s(\mu_R^2)}. \quad (2.9)$$

The parameter μ_R introduces an arbitrary renormalization scale to regulate divergences in the perturbative calculation of α_s . Equation 2.8 shows that the strength of the strong coupling decreases with increasing momentum transfer Q^2 . Therefore, quarks and gluons are *asymptotically free* when probed at high energies. Theoretical work on asymptotic freedom by Gross, Politzer, and Wilczek was rewarded with the 2004 Nobel Prize. On the other hand, as Q^2 approaches Λ , the coupling becomes large and perturbative calculations are no longer possible. This is the regime in which hadronization takes place.

2.2 Cross Sections

In QCD, at high energies and large momentum transfers, interactions between hadrons are due to the hard parton-level process (*hard-scattering*) of the hadron constituents (i.e., quarks and gluons). For example, in the case of $p\bar{p}$ collisions, the hard-scattering process can involve quark-antiquark scattering into a lepton pair, l^+l^- . Equation 2.10 shows the cross section for

such a process. In general, cross sections represent the size of a target measured in units of area (1 barn = 10^{-24} cm²), presented to a probe.

$$\sigma(p(P_1)+\bar{p}(P_2) \rightarrow l^+l^-)+X) = \int dx_1 \int dx_2 \sum_f f_{q/p}(x_1) f_{\bar{q}/\bar{p}}(x_2) \cdot \hat{\sigma}(q(x_1 P_1) + \bar{q}(x_2 P_2)) \rightarrow l^+l^-). \quad (2.10)$$

Here P_1 and P_2 are the proton (p) and anti-proton (\bar{p}) momentum, respectively. $f_{q/p}(x_1)$ and $f_{\bar{q}/\bar{p}}(x_2)$ are *parton distribution functions* (PDFs), which give the probability for a parton q (\bar{q}) to carry a fraction x_1 (x_2) of the proton's (antiproton's) total momentum. $\hat{\sigma}(q(x_1 P_1) + \bar{q}(x_2 P_2)) \rightarrow l^+l^-)$ is the hard-scattering *partonic cross section*. Equation 2.10 separates (*factorizes*) short distance effects, as described by the partonic cross section, from long distance effects, as described by the PDFs. The boundary between these two contributions is defined by the *factorization scale*, μ_F . The factorization scale is not to be confused with the renormalization scale, μ_R , since they do not serve the same purpose. The renormalization scale determines how divergent integrals are replaced with finite expressions, the factorization scale isolates the non-perturbative cross section contributions from the calculable perturbative part.

The PDFs are specific to an initial hadron, i.e. a proton PDF is different from that of a pion. Since PDFs measure properties that cannot be calculated perturbatively, they can only be derived from experiment, usually from *deep inelastic scattering* (DIS) experiments performed at lepton-hadron colliders. PDFs depend on x and Q^2 . Given a PDF at a specific momentum scale, the evolution to any other momentum scale can be determined with the *Dokshitzer-Gribov-Lipatov-Altarelli-Parisi* (DGLAP) equations (45; 46; 47). Since any given experiment




Figure 3. Feynman diagram for LO Z production via $q\bar{q} \rightarrow Z \rightarrow e^+e^-$.

can only cover a limited range of x and Q^2 values, results from different measurements are combined into *global QCD fits*. The PDFs used for this analysis are based on global fits provided by the CTEQ collaboration (Coordinated Theoretical-Experimental Project on QCD).

Figure 3 depicts a graphical representation (*Feynman diagram*) of the hard scattering process for $q\bar{q} \rightarrow Z \rightarrow e^+e^-$. In general, partonic cross sections represented by such diagrams can be derived from the Lagrangian density describing the system under consideration. Every Feynman diagram vertex contributes another factor of the coupling constant to the cross section. The leading order (LO) partonic cross section describing the $q\bar{q} \rightarrow Z$ vertex in Figure 3 is:

$$\hat{\sigma}_{LO}(q\bar{q} \rightarrow Z) = \pi \frac{G_F}{\sqrt{2}} (1 - 4|Q_q|x_W + 8Q_q^2x_W^2) M_Z^2 \delta(\hat{s} - M_Z^2), \quad (2.11)$$

where Q_q is the charge of the quark, and $x_W \equiv \sin^2 \theta_W = 1 - M_W^2/M_Z^2$. G_F is the *Fermi Constant*, defined as $\frac{G_F}{\sqrt{2}} \equiv \frac{4\pi\alpha}{8M_W^2(1-M_W^2/M_Z^2)}$.

2.3 The Measurement of the $Z/\gamma^*(\rightarrow e^+e^-) + \geq n$ Jet Cross Sections

The production of Z/γ^* gauge bosons in association with jets of hadronic particles provides a good opportunity to test perturbative QCD predictions at large momentum transfers. Physical quantities in QCD are generally expressed as perturbative expansions in the strong coupling constant by separating processes at different orders of α_s . Figure 4 and Figure 5 show LO processes for Z/γ^* generation with an additional parton in the final state. The decay of the Z/γ^* gauge bosons into pairs of electrons (e^-) and positrons (e^+) provides an efficient way to identify candidate events with low background contamination.

This analysis studies Z/γ^* gauge boson production and tests the reliability of perturbative QCD predictions over a range of jet energies and jet multiplicities. The results are compared to state-of-the-art MC event simulations, representing LO and next-to-leading order (NLO) QCD calculations.

This measurement is also a significant improvement over previous measurements at the Tevatron (11). The preceding CDF measurement was based on proton-antiproton collisions at a center of mass energy of 1.8 TeV, using $\approx 100 \text{ pb}^{-1}$ of luminosity, and covering jet multiplicities of up to four jets. The measurement presented in this analysis was performed at a higher center of mass energy (1.96 TeV), using $\approx 340 \text{ pb}^{-1}$ of luminosity, and covering jet multiplicities of up to five jets. It is the first measurement at this center of mass energy with the DØ detector.

This analysis also provides a contribution to the study of background processes that are relevant for Higgs boson searches. One of the dominating Higgs production modes at the Tevatron involves the generation of a Higgs boson in association with a Z boson (Figure 6). The




Figure 4. Feynman diagram for Z/γ^* production in association with a parton via $q\bar{q} \rightarrow Z/\gamma^*g$.

study of jet properties and cross sections for processes that generate similar final state particles (Figure 7), results in a more precise distinction between a possible Higgs signal and QCD background.

The analysis was performed with the DØ detector at the Fermilab proton-antiproton Tevatron collider. The following chapter gives an overview of the experimental apparatus that was used for this analysis.




Figure 5. Feynman diagrams for Z/γ^* production in association with a parton via $qg \rightarrow Z/\gamma^* q$.




Figure 6. SM Higgs boson production in association with a Z boson via $q\bar{q} \rightarrow HZ$ (*Higgsstrahlung*).




Figure 7. Feynman diagram for Z/γ^* production in association with two gluons via $q\bar{q} \rightarrow Z/\gamma^* gg$.

CHAPTER 3

EXPERIMENTAL APPARATUS

The Fermi National Accelerator Laboratory (FNAL, or Fermilab) currently operates the world's highest energy proton-antiproton collider, the *Tevatron*. In this chapter the chain of accelerators that is necessary to achieve a center-of-mass collision energy of 1.96 TeV is described. An overview of the DØ detector which is built around one of the collision regions where protons and antiprotons collide is given.

The data used in this analysis was recorded with the DØ detector during the data taking period known as Run II, which officially began in March 2001.

3.1 The Fermilab Accelerators

The Tevatron is the final stage in a sequence of seven accelerators (13; 14; 15). A Cockcroft-Walton *preaccelerator*, a linear accelerator (*Linac*) and a synchrotron (*Booster*) provide a source of 8 GeV protons. The antiproton *Debuncher* and *Accumulator* are two components of the *Antiproton Source*. The *Main Injector* serves as the final boosting stage before injecting protons and antiprotons into the Tevatron. It also provides the necessary source of energetic protons which are needed in the Antiproton Source.

Figure 8 gives an overview of the Fermilab accelerator complex (16).




Figure 8. Schematic view of the Fermilab accelerator chain.

3.1.1 The Preaccelerator

The purpose of the preaccelerator is to produce negatively charged hydrogen ions (H^-) with an energy of 750 keV, which are then transferred into the Linac.

Hydrogen gas (H_2) enters a magnetron surface-plasma source (Figure 9, Reference (13)). Due to the electric field between the anode (negatively charged) and cathode (positively charged) the electrons are stripped away from the hydrogen atoms to create a plasma. The positively charged hydrogen ions then strike the surface of the cathode to collect extra electrons and thereby form negatively charged hydrogen ions. The H^- ions are extracted through the anode aperture with an electric field of 18 kV applied by the extractor plate (see Figure 9).

A commercial Cockcroft-Walton Generator produces a 750 kV potential differential by charging capacitors in parallel from an AC voltage source and discharging them in series, via diodes ¹. The Cockcroft-Walton Generator is used to further accelerate the H^- ions to an energy of 750 keV.

After exiting the Cockcroft-Walton device the H^- ions travel through a transfer line. Before entering into the Linac the continuous stream of H^- ions passes through a single gap radio frequency (RF) cavity which bunches the beam at the RF frequency of the Linac (201.24 MHz).

¹The maximum voltage is limited by how much the air can “stand off” before sparking.




Figure 9. Schematic view of magnetron operation for the hydrogen ion source.

3.1.2 The Linac

The Linac receives bunches of 750 keV H^- ions from the preaccelerator and accelerates them further to an energy of 400 MeV using RF cavities (Figure 10, Reference (13)). The RF cavities are contained within a collection of steel tanks which are holding a sequence of drift tubes separated from each other by gaps. In order to accelerate H^- ions, the cavities are designed in such a way that particles traveling in the gaps experience an acceleration, while particles traveling in the drift tubes are shielded from the RF.

After passing through the Linac, bunches of 400 MeV H^- ions are transferred into the Booster.




Figure 10. Schematic of Linac RF cavity.

3.1.3 The Booster

The Booster is the first synchrotron in the chain of accelerators. It consists of a sequence of dipole and quadrupole magnets and 17 RF cavities arranged in a circle with a diameter of 151 m. The Booster accelerates protons to an energy of 8 GeV.

It is easier to merge negatively charged H^- ions coming from the Linac with protons (H^+ ions) circulating in the Booster due to their opposite charge. Therefore the two beams are merged with the help of dipole magnets, and the electrons are stripped from the H^- ions by letting the combined beam pass through a carbon foil.

Once the Booster is filled with proton bunches, the RF cavities provide an acceleration up to 8 GeV. At the same time the field strength in the dipole magnets is adjusted accordingly in order to maintain a constant radius for the circulating particles. Once the protons have reached an energy of 8 GeV they are transferred into the Main Injector.

3.1.4 The Main Injector

The Main Injector is a circular synchrotron with a diameter of 1 km. It can accelerate both protons (coming from the Booster) and antiprotons (coming from the Antiproton Source) from 8 GeV to 150 GeV, before injecting them into the Tevatron. It also delivers 120 GeV protons to the Antiproton Source.

3.1.5 The Antiproton Source

The Antiproton Source consists of three major components: the *Target Station*, the *Debuncher*, and the *Accumulator*. In the first step the Target Station receives 120 GeV protons from the Main Injector and diverts them onto a Nickel Target. This produces a shower of secondary particles (including antiprotons) at many different angles and large spread in particle momentum. A Lithium lens and bending magnets are used to focus the beam and remove positively charged particles (Figure 11, Reference (17)). A process called *stochastic cooling* is used in both the Debuncher and Accumulator in order to reduce the spread in momentum and position of the antiprotons and thereby “cooling” them.

Both the Debuncher and Accumulator are located in a rounded-triangle shaped tunnel with a circumference of about 51 m. Antiprotons coming from the Target Station are transferred into the Debuncher where the momentum spread of the particles is reduced. It is technically very challenging to accumulate a large quantity of antiprotons. On average for every 1 million protons that hit the Nickel target, only about 20 antiprotons can be gathered. Therefore the Accumulator stores antiprotons until a sufficient amount has been generated that can be




Figure 11. Simplified drawing of anti-proton production with nickel target and lithium lens.

transferred into the Main Injector. The Accumulator must be capable of storing antiprotons over many hours.

3.1.6 The Tevatron

The Tevatron is the final stage in the sequence of proton and antiproton acceleration. It has a diameter of 2 km and uses superconducting magnets which operate at liquid helium temperature providing magnetic fields of up to 4 Tesla. Protons and antiprotons are accelerated to 980 GeV, leading to a center-of-mass collision energy of 1.96 TeV.

Protons and antiprotons travel in groups of particles (*bunches*) in opposite directions while sharing the same beam pipe. A full revolution (*turn*) takes $\approx 21 \mu s$. The Tevatron injects 36

bunches of both protons and antiprotons for each store. A three fold symmetry is imposed by separating the 36 bunches into three superbunches. Overall, this leads to a time structure where bunches of protons and antiprotons (*bunch crossings* or *zero bias events*) collide at 1.7 MHz (18).

3.2 Luminosity and Cross Section

Luminosity \mathcal{L} is a measure of the particle flux per unit area and per unit time ($\text{cm}^{-2}\text{s}^{-1}$). In a collider experiment such as DØ, the luminosity gives an indication of how many proton-antiproton collisions (*events*) occur in a given time and area. The luminosity is determined by measuring the rate of inelastic proton-antiproton scatterings (for each bunch crossing), which have a known cross section (19; 20).

The cross section σ is a measure of the interaction probability per unit flux. Cross sections are usually expressed in *barns*, where $1 \text{ barn} = 10^{-24} \text{ cm}^2$.

The number of times a given process occurs, N , is proportional to \mathcal{L} and σ :

$$N = \sigma \cdot \int \mathcal{L} dt \tag{3.1}$$

$\int \mathcal{L} dt$ is called *integrated luminosity*. Figure 12 shows the integrated luminosity profile of the Tevatron, covering the data-taking period from April 2002 till June 2005. A total integrated luminosity of 343 pb^{-1} was used for the result presented in this analysis.




Figure 12. Tevatron integrated luminosity delivered to DØ (April 2002 - June 2005). The arrow indicates the data taking period during which the data for this analysis was recorded (343 pb^{-1}).

3.3 The DØ Detector

The DØ detector (Figure 13, References (21; 22)) has a magnetic central-tracking system, consisting of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT), both located within a 2 Tesla superconducting solenoidal magnet. Central and forward preshower detectors are located just outside of the superconducting coil. A liquid-argon/uranium calorimeter has a central section (CC) covering pseudorapidities $|\eta|$ up to ≈ 1 , and two end calorimeters (EC) extending coverage to $|\eta| \approx 4$, all three housed in separate cryostats. A muon system resides beyond the calorimetry, and consists of a layer of tracking detectors and scintillation trigger counters before 1.8 Tesla toroids, followed by two more similar layers after the toroids. Luminosity is measured using plastic scintillator arrays located in front of the EC cryostats. The three-tiered trigger and data acquisition systems are designed to accommodate the high luminosities of Run II.

Although a full description of the DØ detector is given in this chapter, the elements that are most relevant for the analysis presented in this dissertation are the calorimeter (Chapter 3.3.4) and tracking system (Chapter 3.3.2).

3.3.1 Coordinate Systems

The coordinate system used for the DØ detector is right-handed. It has the positive z -axis aligned with the direction in which the protons travel and the positive y -axis pointing upwards.

In most cases cylindrical coordinates are used (z, Φ, θ) . Φ is the azimuthal angle in the plane perpendicular to the beam (z -axis), where $\Phi=0$ coincides with the positive x -axis. θ is the polar angle relative to the positive z -axis.




Figure 13. Schematic view of the DØ detector.

Since the colliding protons and antiprotons can exhibit a significant net boost along the z -axis it is suitable to choose a polar angle quantity that is invariant under relativistic transformations. Therefore it is often more convenient to use pseudorapidity η instead of the polar angle θ :

$$\eta = -\ln \cdot \left[\tan \frac{\theta}{2} \right] \quad (3.2)$$

The pseudorapidity approximates the true rapidity,

$$y = \frac{1}{2} \cdot \ln \left[\frac{E + p_z}{E - p_z} \right] \quad (3.3)$$

in the limit of $m \ll E$ (where m is the invariant mass $m^2 = E^2 - p^2$). The term “forward” is used to describe regions at large $|\eta|$ (> 1.1).

In many cases some of the products of a proton-antiproton collision escape along the beam pipe, which makes it difficult to measure momentum components along the z -axis accurately. Therefore it is more convenient to use the momentum vector projected onto a plane perpendicular to the beam axis (*transverse momentum*):

$$p_T = p \cdot \sin \theta. \quad (3.4)$$

In a similar fashion *transverse energy* is defined as

$$E_T = E \cdot \sin \theta. \quad (3.5)$$

3.3.2 Central Tracking System

The central tracking system is essential in measuring the position of the primary interaction vertex with high accuracy, which subsequently allows good measurement of lepton p_T , jet E_T and missing transverse energy (\cancel{E}_T). It can also detect the presence of b -quarks through the measurement of displaced vertices.

The Central Tracking System consists of the silicon microstrip tracker (SMT), the central fiber tracker (CFT) and the superconducting solenoid (Figure 14, Reference (22)). Combining information from both SMT and CFT, the primary vertex resolution is approximately $35 \mu\text{m}$ along the beamline. Jets originating from the decay of b -quarks can be measured with an impact parameter resolution of less than $15 \mu\text{m}$ in $r - \Phi$.

3.3.2.1 Silicon Microstrip Tracker

In order to be able to detect the paths of charged particles emerging from a proton-antiproton collision, the SMT (23) uses wafers of silicon with a thickness of $300 \mu\text{m}$. When a charged particle passes through a positive-negative ($p - n$) junction in silicon it produces electron-hole pairs that can be separated by an applied voltage. The charge which is collected can then be stored in capacitors and later read-out and digitized. The SMT contains of approximately 800,000 individual silicon detectors. Figure 15 shows the design of a basic silicon detector unit (*ladder*) (22).

The SMT is designed in such a way that tracks of charged particles are perpendicular to detector material over a large range of η values. The structure of the device is mostly dictated by the fact that the interaction region is spread out with respect to the center of the detector




Figure 14. The DØ central tracking system with with solenoid, preshower detectors, luminosity monitor, and calorimeter.




Figure 15. Double-sided ladder design, n-side.

($\sigma \approx 25$ cm). This leads to a design of barrel modules combined with disks in the center and larger disks in the forward region (Figure 16, Reference (22)). The SMT has six barrels along the z -axis, each containing four detector layers with a maximal outer radius of 10.5 cm. There are twelve small diameter double-sided “F” disks and four large diameter single-sided “H” disks to cover the far forward region ($|\eta| < 3$). The F-disks are at $|z| = 12.5, 38.2, 43.1$ and 53.1 cm. The centers of the H-disks are located at $|z| = 100.4, 121.0$ cm.

3.3.2.2 Central Fiber Tracker

The Central Fiber Tracker (CFT) (24) is located in between the SMT and the solenoid magnet. The purpose of the CFT is to improve the detection of charged particle tracks within $|\eta| < 2$. It consists of approximately 70,000 scintillating fibers mounted on eight concentric support cylinders with inner and outer radii of 20 and 52 cm, respectively. Each cylinder carries




Figure 16. SMT disk and barrel design.

two layers of fibers running parallel to the beampipe (*axial layers*), and two layers of fibers oriented at small angles of $\pm 3^\circ$ (*stereo layers*) (Figure 17, Reference (25)). The scintillating fibers have a diameter of $835 \mu\text{m}$ and are composed of a scintillating core surrounded by a layer with a high index of refraction, which leads to total internal refraction.

Charged particles passing through a scintillating fiber excite the molecules in the fiber which subsequently release photons in the yellow-green part of the visible light spectrum as they relax to their ground states. Clear fiber waveguides carry the scintillation light to visible light photon counters (VLPCs) that convert the light into electrical signals. The VLPCs are silicon avalanche photodetectors that operate at liquid helium temperature in order to reduce the background due to electronic noise.




Figure 17. a) Location of the Central Fiber Tracker (CFT). b) Closeup view of axial and stereo layers.

3.3.2.3 Solenoidal Magnet

The superconducting solenoidal magnet significantly improves the capabilities of the DØ detector with regard to the detection of charged particle tracks and momentum resolution. The location and physical size of the magnet are constrained by the available space between the inner tracking system and the vacuum vessel of the central calorimeter. The magnet has a length of 2.73 m and a diameter of 1.42 m and provides uniform field of 2 T (20 kG) over most of the region covered by the inner tracking system (Figure 18, Reference (22)). Figure 19 shows a perspective view of the solenoid inside the central calorimeter with its chimney and control dewar (22).

3.3.3 Preshower Detectors

The preshower detectors (26; 27) are designed to improve the identification of electrons and background rejection. By the time a charged particle has reached the preshower detectors it has already passed through several detectors as well as other upstream material like cables and support structures. The preshower detectors allow to correct for upstream energy losses in the offline event reconstruction. Due to their fast response time the preshower detectors can also be used for Level 1 triggering (see Chapter 4.1).

Scintillators are used to detect both position and energy of charged particles. In contrast to the scintillators used in the CFT, preshower scintillators are triangular shaped (Figure 20, Reference (22)). This allows to arrange layers of scintillators without creating any dead space and thereby improving the accuracy of position measurements. The center of each scintillator




Figure 18. $y - z$ view of the DØ magnetic field with both the toroid and solenoid magnets at full current. Numbers are in kG (10 kG = 1 T).

carries a wavelength-shifting fiber which collects the light created by passing charged particles.

The light is transmitted via clear fibers to VLPCs for readout.

3.3.3.1 Central Preshower Detector

The Central Preshower Detector (CPS) is located in the 5 cm gap between the solenoid and the central calorimeter, covering the region $|\eta| < 1.3$ (Figure 14, Reference (22)). It consists of a layer of lead radiator which has a thickness corresponding to approximately one radiation-length (X_0), followed by three layers of triangular scintillator strips. The purpose of the lead radiator is to ensure that the total amount of material encountered by a passing particle is




Figure 19. Perspective view of the solenoid inside the central calorimeter.




Figure 20. Cross section and layout geometry of the CPS and FPS scintillator strips.

constant at all pseudorapidities. The scintillating layers are arranged in an axial- u - v geometry, with a u stereo angle of 23.8° and a v stereo angle of 24.0° . Each layer has a total number of 2,560 readout channels.

3.3.3.2 Forward Preshower Detector

The two Forward Preshower Detectors (FPS) are attached to the faces of the end calorimeters and cover a region of $1.5 < |\eta| < 2.5$ (Figure 14, Reference (22)). Each detector consists of an upstream double layer of scintillator strips (minimum ionizing particle layers, or MIP layers), followed by a lead-stainless-steel absorber layer, and another double layer of scintillator strips behind it (shower layers). This configuration allows for the distinction between electron-like

and photon-like objects. While an electron-like object will create a minimum ionizing energy deposit in the MIP layers, a photon-like object will not. Therefore electrons can be identified by requiring spatially matched energy deposits in both the MIP layers and the shower layers, whereas photons will only have energy deposits in the shower layers.

3.3.4 Calorimeter

The main purpose of the calorimeter system (28) is to measure the energy deposits from electrons, photons, and jets. It also allows direct identification of electrons, photons, jets, and muons. In addition, measurements of the transverse energy balance in an event can detect the presence of neutrinos.

The calorimeter system consists of a central calorimeter (CC) covering $|\eta| < 1.2$ and two end calorimeters (EC), covering $1.3 < |\eta| < 4.0$ (Figure 21, Reference (22)). Each of the calorimeters has an electromagnetic section, followed by fine and coarse hadronic sections (FH and CH respectively). Since liquid argon is used as the active medium, all calorimeters are contained within cryostats. Different types of materials are used for absorber plates:

- 3 mm (4 mm) plates of depleted uranium for the CC (EC) electromagnetic sections.
- 6 mm plates of uranium-nobium (2%) for the fine hadronic sections.
- 46.5 mm plates of copper (stainless steel) for the CC (EC) coarse hadronic sections.

A typical calorimeter cell is shown in (Figure 22, Reference (25)). Each cell consists of a grounded absorber plate and a signal board maintained at a positive high voltage of typically 2 kV. The 2.3 mm gap between the absorber plate and signal board is filled with liquid argon.

The calorimeter cells are arranged to form pseudo-projective towers (Figure 23, Reference (22)).

In order to measure the energy of electromagnetically interacting objects, the calorimeter takes advantage of the electromagnetic shower process. For example, an incoming electron will emit Bremsstrahlung photons when passing through the dense absorber material and will interact with the nuclei in the absorber via Coulomb interaction. The emitted photons will subsequently decay into electron-positron pairs. This chain of processes continues until the energies of the secondary particles fall below the threshold for pair production. After that the particles mainly interact via ionization. At each stage of the electromagnetic shower, charged particles are ionizing the liquid argon. The high voltage between the absorber plates and signal boards is then used to collect the ionization charges, which allows a measurement of the energy in the shower. Typical transverse sizes of electromagnetic showers are in the range of 1-2 cm.

Hadronic showers are induced by the interaction between hadronic particles and the nuclei of the absorber material via the strong nuclear force. Secondary hadronic particles then further interact via inelastic nucleus collisions till their energy falls below a threshold. Typical transverse sizes of hadronic showers are of the order of 10 cm. The space in between the central and end calorimeters ($1.1 < |\eta| < 1.4$) is referred to as the intercryostat region (ICR). In order to be able to measure the energies of particles that pass through this gap in the calorimeter coverage, additional detectors are used. Calorimeter cells called massless gaps (MG) are installed before the first layer of uranium inside of the central and end cryostats. Additionally, a ring of scintillator tiles mounted to the exterior surface of the end cryostats comprises the intercryostat detector (ICD).




Figure 21. Isometric view of the central and two end calorimeters.

Figure 24 shows the different calorimeter detection layers for a given η value (29).

3.3.5 Muon System

Due to their large mass and long lifetime, muons pass through the calorimeter by depositing only a small amount of energy (≈ 2.5 GeV) via ionization. Therefore, the outermost subdetector is dedicated to the detection of muons (Figure 25, Reference (25)). The muon system is separated into central and forward detectors. A 1.9 T iron toroid magnet is used for muon momentum measurements.

Proportional Drift Tubes (PDT), Mini Drift Tubes (MDT), and scintillators are the main




Figure 22. Schematic view of two calorimeter cells.

detection elements used in the muon system. Drift tubes collect the ionization charges created by muons passing through a gas mixture onto high voltage wires. Correlating the arrival times of ionization charges from different drift tubes with the beam crossing time, allows to extrapolate the path of muons as they pass through the detector. Scintillators are mainly used for their good timing resolution (≈ 4 ns) which allows a way to trigger on muons.

The central muon system (30) covers the region of $|\eta| < 1.0$ and is referred to as the Wide Angle Muon System (WAMUS). It consists of three PDT layers, with the first layer (A layer) in between the toroid magnet and the calorimeter, and two more layers (B and C layers) after the toroid magnet. Additional layers of scintillators before the A layer and covering the outside of the muon system allow for spatial and precise timing measurements to reject cosmic rays.




Figure 23. Schematic view showing the calorimeter segmentation pattern. The shading pattern indicates cells for signal readout. The radial lines show the detector pseudo-rapidity intervals.




Figure 24. Schematic view of different calorimeter detection layers vs η .




Figure 25. The DØ muon system.

The forward muon system (31) covers the region of $1.0 < |\eta| < 2.0$ and is referred to as the Forward Angle Muon System (FAMUS). It consists of three MDT layers and scintillators, with the first layer (A layer) before the toroid magnet, and two more layers (B and C layers) after the toroid magnet.

CHAPTER 4

THE DØ TRIGGER AND DATA ACQUISITION SYSTEMS

DØ uses a three level trigger system (25; 32) to handle the high proton-antiproton collision rate of 1.7 MHz. Each succeeding level of triggering processes fewer events, with more sophisticated trigger algorithms. At the first stage the Level 1 (L1) system uses a hardware trigger to reduce the event rate to ≈ 1.5 kHz. At the next stage the Level 2 (L2) system further reduces the event rate to ≈ 800 Hz. L2 uses hardware engines associated with specific detector subsystems and a single global processor for the final L2 trigger decision. In the last step the Level 3 (L3) system which consists of a farm of microprocessors reduces the event rate to ≈ 50 Hz. Only those events that pass all three trigger levels are stored for further offline reconstruction and analysis.

4.1 The Level 1 Trigger

The L1 trigger system uses information from the tracking, preshower, calorimeter, and muon subdetectors to provide an event rate reduction by a factor of ≈ 1000 . Field programmable gate arrays (FPGAs) check whether a given event satisfies one of up to 128 L1 trigger conditions. If at least one trigger condition is satisfied the event information is digitized and buffered to await a L2 decision. L1 trigger decisions are made within a $3.5 \mu\text{s}$ time window.

4.1.1 The Level 1 Calorimeter Trigger

For trigger purposes the calorimeter is segmented into 1280 trigger towers of $\Delta\eta \times \Delta\Phi = 0.2 \times 0.2$. Level 1 calorimeter (L1CAL) trigger decisions are based on the amount of transverse energy deposited in the electromagnetic layers (EM) and electromagnetic plus fine hadronic layers (TOT) of the trigger towers. L1CAL trigger conditions require that a specific number of EM or TOT trigger towers be above a certain transverse energy threshold. In addition L1CAL can also impose thresholds on the total sum of transverse energy and the missing transverse energy in a given event.

4.1.2 The Level 1 Central Track Trigger

The Level 1 Central Track Trigger (L1CTT) uses information from the CFT and CPS sub-detectors in the central region ($|\eta| < 1.7$), and information from the CFT and FPS subdetectors in the forward region ($1.4 < |\eta| < 2.5$). Possible track candidates are identified by FPGAs using hit patterns in the axial layers of the CFT and matching energy deposits in the preshower detectors (Figure 26, Reference (22)). The p_T of a track candidate can be estimated by the azimuthal bend of the CFT hits. L1CTT track candidates are assigned into p_T bins according to their transverse momenta: 1.5-3 GeV; 3-5 GeV; 5-10 GeV; or above 10 GeV. Trigger conditions can be specified by a certain number of tracks above a p_T threshold, with or without the requirement of a CPS cluster match to the track.

4.1.3 The Level 1 Muon Trigger

The Level 1 Muon Trigger (L1Muon) uses information from the muon wire chambers, muon scintillators, and L1CTT tracks. L1Muon trigger logic is implemented in FPGAs and trigger




Figure 26. A hypothetical L1CTT track with hits in eight CFT axial doublet layers and the CPS axial layer.

conditions require a combination of criteria based on p_T thresholds, geographical region, track quality, and multiplicity.

4.2 The Level 2 Trigger

The L2 trigger system is designed to reduce the L1 event rate by a factor of <10 . It receives inputs from both the L1 system and the detector subsystems. L2 operates in two stages. In the first stage, subdetector-specific preprocessors form physics objects such as electrons, jets, or tracks. Individual preprocessors for the tracking, preshower, calorimeter, and muon subdetectors run in parallel and are located in separate crates. L2 preprocessor physics objects

are then used at the second stage by a global processor. The L2 global processor (L2GBL) makes the final L2 trigger decision by imposing selection criteria on the preprocessor physics object, including correlations between multiple detector subsystems.

The two-stage L2 architecture (*stochastic pipeline*) can make trigger decisions within a 100 μs time window (see Appendix B for a detailed description of the L2 trigger system). Events passing L2 trigger requirements are flagged for full detector readout and further refined analysis at the L3 triggering stage.

4.2.1 The Level 2 Calorimeter Preprocessor

The Level 2 calorimeter preprocessor (L2Cal) receives the full list of 2560 EM and TOT trigger towers from L1CAL.

The L2 jet algorithm forms jet objects by clustering 5×5 groups of TOT trigger towers centered around seed towers. A jet seed tower is any TOT trigger tower with $E_T \geq 2 \text{ GeV}$. The list of seed towers for the L2 jet algorithm is E_T -ordered. Depending on the configuration, overlapping L2 jet candidates can be reported as separate jets or simply as the highest- E_T jet. An E_T -ordered list of all L2 jets is sent to L2GBL.

The L2 Electron/photon algorithm forms EM objects by clustering the transverse energies of EM seed towers with their largest E_T neighbors. An EM seed tower is any EM trigger tower with $E_T \geq 1 \text{ GeV}$. The list of seed towers for the L2 EM algorithm is E_T -ordered. The algorithm also reports a list of EM objects based on information from seeds only (without considering the neighboring towers). To reduce background the EM energy fractions for both the leading and next-to-leading EM cluster that form the EM object are calculated. Additionally, the isolation

of the EM cluster from other energy deposits in the calorimeter is evaluated. An E_T -ordered list of all L2 EM objects and parameters is sent to L2GBL.

4.2.2 The Level 2 Muon Preprocessors

The Level 2 Muon system (L2Muon) consists of two components: a preprocessor for the central muon region (L2 Muon Central, or L2MUC) and a preprocessor for the forward muon region (L2 Muon Forward, or L2MUF). L2Muon receives inputs from L1 (L1Muon) and the muon subdetector drift chambers and scintillators. Inputs to L2Muon are first received by *second level input computers* (SLICs), where most of the processing is done. A list of muon candidates with each containing information about η , ϕ , p_T , sign, and timing is sent to L2GBL for further processing.

4.2.3 The Level 2 Preshower Preprocessor

The Level 2 preshower preprocessor (L2PS) receives inputs from the L1 trigger system. Based on preshower detector information it forms clusters of energy deposits with η and ϕ coordinates that can then be matched with calorimeter trigger towers or CFT trigger tracks at the L2GBL processing stage.

4.2.4 The Level 2 Tracking Preprocessor

The Level 2 tracking preprocessor uses information from L1CTT and the SMT to form lists of L2 track candidates that can be used by L2GBL for triggering. It triggers on vertices that are displaced from the primary interaction vertex. Displaced vertices are characteristic of long-lived particles such as B -Mesons, and can therefore be used to identify heavy-flavored quarks.




Figure 27. The definition of roads based on L1 tracks and SMT hit selection in L2STT.

The L2 tracking preprocessor can be operated in two different modes. In the first mode of operation (L2 Central Track Trigger, or L2CTT) it further refines tracking information coming from L1CTT. In the second mode of operation (L2 Silicon Track Trigger, or L2STT) it combines L1CTT and SMT information. Figure 27 shows the conceptual design of L2STT (22). It uses hit information from the first and last layer of the CFT to define a *road*, and then extrapolates that road into the SMT to find additional hits in the axial strips of the SMT silicon ladders.

4.2.5 The Level 2 Global Processor

The Level 2 global processor (L2GBL) receives lists of trigger objects that are generated by the L2 preprocessors (L2Cal, L2MUC, L2MUF, L2PS, L2CTT, L2STT). It creates global trigger

objects by either using the trigger objects generated by the preprocessors or by combining trigger object information from different preprocessors. For example, spacial correlations between track candidates and EM objects in the calorimeter can be used to select a cleaner sample of electron candidates. L2GBL makes the final L2 trigger decision by imposing cuts on global trigger objects that are defined by trigger list information which is parsed from the trigger control computer.

4.3 The Level 3 Trigger and Data Acquisition

The L1 and L2 trigger systems do not use the full detector readout for their trigger decisions. The event rate needs to be reduced to 1 kHz or below, so that the Level 3 (L3) trigger system can take advantage of the full detector readout. L3 then further reduces the event rate to ≈ 50 Hz. The average event size is ≈ 200 kBytes.

The L3 trigger system is based on sophisticated reconstruction algorithms that resemble the algorithms for offline event processing (see Chapter 5) as closely as possible, given restrictions due to available processing power.

The data acquisition system (L3DAQ) is based on a single Cisco 6509 Ethernet switch that transfers data at a rate of < 250 Mbyte/s. Detector data is transferred from commodity VME single-board computers (SBCs), via the Ethernet switch to the individual L3 farm nodes. A supervisor process running on a separate CPU provides the interface between the main DØ run control system (COOR) and L3DAQ.

After passing L3 trigger requirements the data are buffered to local disks and finally transferred to a permanent storage facility.

CHAPTER 5

OFFLINE EVENT RECONSTRUCTION

This chapter describes how candidate electrons, jets, tracks, and vertices are *reconstructed* from the raw detector data. A collection of complex software algorithms written in C++ called *d0reco* (33) is used for this reconstruction process. As described in the previous chapter, a stream of digital readout signals from the subdetectors is recorded for each event that passes all three levels of triggering. Powerful PC computing farms are deployed to analyze the data and reconstruct candidates of physical objects (electrons, jets, tracks, etc) that can then be used in the final analysis.

5.1 Track Reconstruction

In order to reconstruct track candidates, two different types of algorithms are used. The *Alternative Algorithm* (AA) (34) which uses a road-following method, and the *Histogramming Track Finder* (HTF) (35) which relies on a histogramming method.

The AA method starts from any combination of three hits in the SMT barrels or disks. Moving outwards, towards the CFT, it then extrapolates the sequence of hits to the next SMT or CFT layer. If a hit is found within a search window, a χ^2 test is performed. If the χ^2 value is below a certain threshold, the newly found hit is associated with the track candidate. If no hit is found, it is recorded as a *miss*. Construction of track candidates ends when the last layer of the CFT is reached, or when three misses are recorded.

The HTF method takes advantage of the fact that the trajectory of a charged particle moving perpendicular to a homogeneous magnetic field can be characterized by three parameters: ρ , d_0 , and Φ , where ρ is the radius of curvature, d_0 is the distance of closest approach with respect to (0,0) (*impact parameter*), and Φ is the direction of the track at the point of closest approach to (0,0). For track candidates with small impact parameters, every pair of hits in (x,y) coordinate space that belongs to the same track corresponds to a single point in (ρ, Φ) parameter space. Therefore, by examining every pair of hits and filling a 2-dimensional (ρ, Φ) histogram, a peak in the histogram would correspond to a track candidate.

Both track reconstruction algorithms generate lists of track candidates. A final list of tracks is generated after eliminating duplicates. The tracks in the final list are sorted by number of hits, fewest misses, and lowest χ^2 value. The best track is automatically kept, and the rest of the tracks are examined based on these parameters. Figure 28 shows an example of hits and reconstructed tracks (36).

5.2 Primary Vertex Reconstruction

The primary vertex (PV) is the location of the proton-antiproton hard scatter collision. It is important to identify the position of the PV with high accuracy, since it is an essential ingredient to reconstruct jets, electrons and missing transverse energy. It is also important to select an algorithm (37) which distinguishes between the hard scatter vertex and the vertices from additional *minimum bias* (MB) interactions.

Based on the list of reconstructed tracks (see Section 5.1), a list of vertices is generated by extrapolating the tracks back to the z -axis. Clusters of vertices are then formed by selecting




Figure 28. Axial view (looking down the beam-pipe) of a recorded event showing hits and reconstructed tracks. CFT hits are represented by squares, and SMT hits are represented by circles. Hits are colored solid if they are associated with a reconstructed track (solid lines). The curvature of the reconstructed tracks is due to the solenoidal magnetic field, which is pointing out of the page.

vertices within 2 cm of each other along the z -axis. In each vertex cluster, the vertex with the highest number of tracks pointing at it is stored in a list of “selected” vertices. For every selected vertex, nearby tracks are used to compute the probability that the vertex does not come from a MB interaction. The computation of this probability is based on the assumption that tracks coming from MB interactions will have a smaller transverse momentum compared to tracks coming from the hard scatter. Finally, the vertex with the smallest MB probability is chosen as the PV.

5.3 Electromagnetic Object Reconstruction and Identification

Electromagnetic objects (EM objects), such as electrons and photons, are initially identified based on calorimeter information. Since photons do not leave signals in the tracking system, a track matched to the energy deposit in the calorimeter distinguishes electrons from photons.

EM object reconstruction begins with forming initial calorimeter clusters. Different algorithms can be used to find those initial clusters:

- Simple-Cone tower clustering algorithm (“Scone Method”)
- Cell Nearest Neighbor clustering algorithm (“CellNN Method”)
- Track extrapolation clustering algorithm (“Road Method”)

In this analysis, objects reconstructed with the simple cone algorithm are used.

5.3.1 Simple-Cone Clustering Algorithm

The simple-cone algorithm (38) clusters calorimeter cells based on precision readout data around seeds with $E_T > 1.5$ GeV in a cone of radius $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} < 0.2$. Additionally, for each cell the ratio of the energy in the EM layers to the total energy (based on CPS, EM, and FH layers) is required to be above 0.9¹. Next, all clusters satisfying the above criteria are tested for *isolation*:

$$Isolation = \frac{E_{tot}(R < 0.4) - E_{EM}(R < 0.2)}{E_{EM}(R < 0.2)}, \quad (5.1)$$

where $E_{EM}(R < 0.2)$ is the EM energy within a cone of radius $R < 0.2$ (based on EM layers), and $E_{tot}(R < 0.4)$ is the total energy within a cone of radius $R < 0.4$ (based on EM, FH, and CH layers). All initial EM clusters are required to have an isolation of less than 0.2. The isolation parameter gives a measure of how deep and narrow a given cluster is. EM objects tend to deposit most of their energy in a narrow region of the EM layers, while hadrons deposit their energies in the hadronic layers in a much wider radius.

5.3.2 Electromagnetic Object Identification Parameters

Various parameters are calculated for every EM cluster that is formed by the simple-cone algorithm. This gives flexibility when defining EM objects at the analysis stage.

ID: All EM clusters are assigned an ID of 10. If in addition a cluster has a track loosely matched (in η and Φ) to it, it is assigned an ID of ± 11 (“+” for electrons, “−” for positrons).

¹This definition of EM fraction is different from the definition that is used offline (see Section 5.3.2).

Isolation: An isolation cut of 0.2 is already applied at the reconstruction stage (see Section 5.3.1).

Electromagnetic Fraction: The electromagnetic fraction (EM fraction) discriminates between EM and hadronic calorimeter energy deposits. It takes advantage of the fact that EM showers are almost entirely contained within the EM layers of the calorimeters. EM fraction is defined as:

$$EMfraction = \frac{E_{EM}(R < 0.2)}{E_{tot}(R < 0.2)}, \quad (5.2)$$

where $E_{EM}(R < 0.2)$ is the EM energy within a cone of radius $R < 0.2$ (based on EM layers), and $E_{tot}(R < 0.2)$ is the total energy within a cone of radius $R < 0.2$ (based on EM, FH, and CH layers).

H-Matrix: The H-Matrix distinguishes between EM and hadronic energy deposits, by analyzing the longitudinal and transverse shape of the showers. Based on MC generated electrons a covariance matrix (M) is defined using a set of seven discriminant variables :

$$M_{ij} = \frac{1}{N} \sum_{n=1}^N (x_i^n - \langle x_i \rangle) (x_j^n - \langle x_j \rangle), \quad (5.3)$$

where x_i^n is the value of variable i for electron n , and $\langle x_i \rangle$ is the mean value of variable i . The seven variables that are used are listed below:

- Shower energy fraction in 1st, 2nd, 3rd, and 4th EM layer of the calorimeter.
- Cluster size in $r - \Phi$ based on the 3rd EM layer of the calorimeter.

- Total shower energy.
- Primary vertex position.

The H matrix is defined as the inverse of the covariance matrix (Equation 5.3):

$$H \equiv M^{-1} \quad (5.4)$$

Using the H matrix one can calculate a χ^2 -like variable that gives a measure of the likelihood that a given shower k is consistent with an EM object shower:

$$\chi^2 = \sum_{ij} (x_i^k - \langle x_i \rangle) H_{ij} (x_j^k - \langle x_j \rangle) \quad (5.5)$$

Track Matching: Requiring that a track is associated with a calorimeter EM cluster is a powerful discriminant between electrons and photons. In this analysis *global tracks* are used for tracking confirmation, i.e. tracks based on information from both the CFT and SMT subdetectors. Using calorimeter and tracking information the following χ^2 variable is calculated:

$$\chi^2 = \left(\frac{\Delta\Phi}{\sigma_\Phi} \right)^2 + \left(\frac{\Delta z}{\sigma_z} \right)^2 + \left(\frac{E_T/p_T - 1}{\sigma_{E/p}} \right)^2 \quad (5.6)$$

where in Equation 5.6:

- $\Delta\Phi$ (Δz) is the difference in Φ (z) between the EM cluster position in the 3rd EM calorimeter layer and the extrapolation of the track to the same layer¹.
- σ_Φ , σ_z , and $\sigma_{E/p}$ are the root-mean-squares of the experimental measurements of each quantity.
- E_T/p_T is the ratio of the transverse energy of the EM calorimeter cluster and the transverse momentum of the track.

A track is matched to an EM cluster by requiring that the track matching χ^2 probability is $\chi^2 > 10^{-2}$.

5.4 Jet Reconstruction and Identification

Individual quarks and gluons cannot be detected directly due to color confinement. Instead, collimated streams of hadrons (*jets*) are observed in the detector. In this section the techniques that are used to reconstruct and identify jets are discussed.

5.4.1 Jet Cone Algorithm

The *Run II Midpoint Jet Cone Algorithm* is used to reconstruct jets for this analysis (39; 40). The general idea behind a jet cone algorithm is to define a cone in $\eta \times \Phi$ space, and to group together particles whose trajectories lie within that cone. After choosing an initial trial axis for a cone, the energy-weighted centroid is calculated based on contributions from all particles within the cone. Using the new cone axis, this process is iterated until a “stable” position is

¹EM showers typically deposit the bulk of their energy in the 3rd EM layer.

found, where the centroid of the energy depositions within the cone is aligned with the geometric axis of the cone.

Input into the algorithm can be based on calorimeter towers, in case the jet reconstruction is performed at the detector level in data or MC. Hadrons (partons) can be used as input at the particle (parton) level in MC.

The following is a description of the Run II Midpoint Jet Cone Algorithm:

1. Only *seed particles* with $p_T^{seed} > 0.5$ GeV are considered for jet reconstruction.
2. Cones are formed based on 4-vector variables (*E-scheme*):

$$p = (E, \mathbf{p}) = \sum_i (E^i, p_x^i, p_y^i, p_z^i), \quad (5.7)$$

$$p_T = \sqrt{(p_x)^2 + (p_y)^2}, \quad (5.8)$$

$$y = \frac{1}{2} \ln \frac{E + p_z}{E - p_z}, \quad (5.9)$$

$$\Phi = \tan^{-1} \frac{p_y}{p_x} \quad (5.10)$$

where p is the energy-momentum 4-vector of the cone, and the sum runs over all particles i within a cone of radius 0.5. p , y , and Φ define the centroid for a given cone. Each stable cone is added to a list of *proto-jets*. Only proto-jets with $p_T > 8.0$ GeV are considered.

3. Midpoints between any combination of two proto-jets are also used as seeds and iterated until stable cones are found. This is done to remove sensitivity to soft radiation (Figure 29, Reference (39)).




Figure 29. Illustration showing how the presence of soft radiation between two jets may cause a merging of the jets (right) that would not occur in the absence of the soft radiation (left).

4. The proto-jets are checked for overlapping regions. In order to avoid double counting of energy a merging/splitting algorithm is applied. Two proto-jets are *merged* if the shared energy between them is greater than half of the lower p_T proto-jet. Otherwise, the two proto-jets are *split* and the tower energies are assigned to the closest proto-jet.

5.4.2 Jet Identification Parameters

A set of quality cuts is applied to every reconstructed jet in order to reduce fake jets from calorimeter noise.

EM Fraction (EMF): Hadronic shower formation tends to deposit a significant energy fraction in the hadronic layer of the calorimeter, whereas EM objects mostly shower in the EM layers. Therefore, a cut on the fraction of transverse energy in the EM layers distinguishes jets from EM objects.

Coarse Hadronic Fraction (CHF): Coarse hadronic fraction is the fraction of transverse momentum of a jet that is deposited in the coarse hadronic layers of the calorimeter.

Hot Fraction (HotF): Hot Fraction is the ratio of transverse energy in the most energetic tower to that of the next most energetic tower in the jet. If a significant amount of the total jet energy is originating from a single calorimeter tower, it is likely that the tower is generating artificially high read-out signals due to detector problems (*hot tower*). A cut on the Hot Fraction parameter eliminates jets originating from hot towers.

N90: N90 is the number of towers making up 90% of the jet energy. A cut on the N90 parameter eliminates jets originating from hot towers as well.

L1 Confirmation: L1 confirmation was introduced in order to deal with precision readout noise problems. The jet energy at the L1 trigger tower level is compared with the jet energy derived from the jet cone algorithm, which is based on calorimeter cell precision readout:

$$L1conf = \frac{\sum_{trigger} E_T^i}{p_T^{jet} \cdot (1 - CHF)} \quad (5.11)$$

where $\sum_{trigger} E_T^i$ is the sum of transverse TOT trigger tower energies (with respect to the center of the detector) within $\Delta R < 0.5$ of the reconstructed jet; p_T^{jet} is the transverse momentum of the reconstructed jet (with respect to the reconstructed primary vertex); and CHF is the Coarse Hadronic Fraction of the reconstructed jet.

5.4.3 Jet Energy Scale

The jet energy scale (JES) calibrates the reconstructed energy of the jets to the energy of the jets at the particle (or hadron) level, i.e. before the particles of the jet hit the detector (41). Depending on the jet p_T and jet η , the JES applies a calibration factor to obtain the particle level jet energy ($E_{jet}^{particle}$) from the measured jet energy ($E_{jet}^{calorimeter}$) according to the following relation:

$$E_{jet}^{particle} = \frac{E_{jet}^{calorimeter} - E_{offset}}{R_{jet} \cdot R_{cone}} \quad (5.12)$$

where E_{offset} is the *offset energy* within a jet, R_{jet} is the *jet response* correction, and R_{cone} is the *out of cone showering* correction.

Offset Energy: Energy contributions that are not related to the physics processes that are responsible for creating a jet are subtracted from the measured jet energy. The offset energy term contains contributions from multiple interactions, underlying event energy, electronic noise, uranium noise, and pile-up from previous bunch crossings.

Jet Response: This contribution measures the calorimeter response of the hadronic particles and the amount of energy within a jet that is lost due to uninstrumented regions and dead material in the calorimeter.

Out of Cone Showering: This parameter takes into account the fraction of the particle jet energy that is deposited inside the algorithm cone. The jet energy is corrected back to the particle level, and therefore calorimeter showering effects must be removed. R_{cone} is




Figure 30. The jet energy scale correction factor measured for jets in data as a function of E (top) and η (bottom).

less than one, since the effect of showering is a net flux of energy from the inside to the outside of the cone.

Figure 30 shows the overall jet energy scale corrections as a function of E and η .

CHAPTER 6

MONTE CARLO EVENT SIMULATION

MC event simulation programs are an essential tool in particle physics (42; 43). The simulations represent the theoretical understanding of nature, and the ultimate goal is to simulate nature as accurately as possible. In particle physics, MC simulations aim to include all stages of interactions that occur at high energy collisions including initial creation of partons in hadronic collisions (*parton level*), subsequent hadronization (*particle level*), and simulations of detector responses (*detector level*). Therefore, MC event simulations compare MC output with data in order to test theoretical predictions. The simulations can also be used to estimate detector acceptances and to study background processes. Alternatively, the simulations permit to correct data for detector effects so that direct comparisons to generator level MC simulations can be made.

This chapter gives a description of the MC simulation tools and techniques, that were used in this analysis.

6.1 The PYTHIA Event Generator

PYTHIA is a general purpose MC event generator (44) which describes most physical aspects of a typical high-energy event:

1. The initial beam particles, such as protons and antiprotons, are characterized by PDFs.

2. One parton from each initial particle starts to radiate through a *Bremsstrahlung*-like process, such as $q \rightarrow qg$, to initiate a sequence of branchings which builds up an initial-state shower (*Initial State Radiation*, or ISR).
3. Two partons participate in the *hard process* which produces a number of outgoing particles. PYTHIA is optimized for hard processes that have two particles in the initial and final states, such as $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$ ($2 \rightarrow 2$ process). Intermediate resonances, like Z/γ^* , need to be included in the description of the hard process.
4. Final state particles can also radiate to initiate a final-state shower (*Final State Radiation*, or FSR).
5. The remaining partons (*spectators*) in the two incoming hadrons may interact.
6. Beam remnant interactions are taken into account.
7. Outgoing quarks and gluons are forming color neutral hadrons due to QCD confinement.
8. Many of the produced hadrons are unstable and decay further.

In order to describe inclusive $Z/\gamma^* \rightarrow e^+e^-$ events, a PYTHIA sample based on the $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$ hard process is used, where the LO cross section is due to the electroweak interaction. ISR parton showers provide QCD corrections to the LO calculation. Figure 31 shows an illustration of a parton shower (9). Based on the DGLAP equations, which give the probability for a parton to radiate, or branch, the parton shower is evolved from the scale Q down to a cut-off scale Q_0 where it terminates. At this scale, α_s becomes large and perturbation theory breaks down. At that stage the colored partons are converted into colorless hadrons

by the hadronization process. PYTHIA uses a phenomenological model known as the *String Model* to describe this process. As a color-confined quark-antiquark pair moves out in opposite directions, the color field between them collapses into a string-like configuration, where each string has a uniform energy per unit length, and the potential energy in the string increases as the quarks move apart. Once the potential becomes energetically favorable another $q\bar{q}$ pair is pulled from the vacuum, and the string split into two. This process continues until only on mass-shell hadrons remain.

6.2 Combining Matrix Elements with Showering

PYTHIA produces multijet events by incorporating $2 \rightarrow 2$ hard processes with parton-showering simulation of QCD radiation that is subsequently interfaced with the string hadronization model. Since parton showers are only approximations of soft parton radiation in the limits of low p_T and small emission angles (*soft* and *collinear limits*), final states with several high p_T and well separated jets are not properly described. Such final states occur frequently at the Tevatron due to the large momentum transfer, Q^2 . In order to give a more accurate description of such processes, matrix element generators are combined with parton showering simulations by separating phase space into two regions, separated by a p_T *matching threshold*. In the high p_T regime, jet production is modeled by providing the appropriate matrix elements; the low p_T region is described by the parton showering simulation. Exact book-keeping is necessary when combining the two regions in order to avoid the double counting of parton configurations (48; 49).




Figure 31. Schematic view of a parton shower.

6.2.1 Combining ALPGEN with PYTHIA

In order to describe inclusive $Z/\gamma^* \rightarrow e^+e^- + \geq n$ jet events ($n \geq 1$), the tree-level matrix element generator ALPGEN (51; 52) is used. ALPGEN was designed to simulate specific SM processes in hadronic collisions, emphasizing large jet multiplicities in the final state. ALPGEN calculates parton level final states to lowest order in perturbation theory. The ALPGEN output is interfaced with PYTHIA for showering and hadronization.

No procedure to avoid double counting of parton configurations is implemented in the ALPGEN + PYTHIA samples used for this analysis.

6.2.2 Combining MADGRAPH with PYTHIA

MADGRAPH (53) is a tree-level matrix element generator, and it is based on specifying initial and final state particles for any tree level SM process. It then creates a list of all relevant Feynman diagrams and calculates the corresponding matrix elements. The program is able to calculate matrix elements for any SM process. The only limitation is processing power¹. The MADGRAPH output is interfaced with PYTHIA for showering and hadronization.

The matching between MADGRAPH and PYTHIA in order to avoid double counting is done following a modified CKKW prescription (54; 55; 50). These samples are referred to as *CKKW samples* and are used for the final cross section comparison.

¹At present MADGRAPH is limited to 10,000 diagrams per subprocess, corresponding to $W + 5$ jets.

6.3 The MCFM Monte Carlo Simulation

MCFM (*Monte Carlo for FeMtobarn processes*) (56; 57; 58) can calculate parton level cross sections for $Z/\gamma^*(\rightarrow e^+e^-)$ at NLO in α_s for up to two partons in the final state.

The final analysis results are compared to MCFM cross sections.

6.4 The DØ Detector Simulation

After all final state particles are generated, detector effects can be simulated with a simulation package called D0GSTAR (*D0 GEANT Simulation of the Total Apparatus Response*) (59). D0GSTAR is based on the CERN package GEANT (*GEometry ANd Tracking*) (60) which simulates the passage of particles through matter. Effects such as ionization, showering, and magnetic field interaction in the DØ detector are modeled. The output of this simulation has the same format as the real data recorded by the detector. The full event reconstruction applied to real data can also be applied to the simulation output for direct comparisons.

CHAPTER 7

DATA AND MONTE CARLO SAMPLES

This chapter gives a description of the data and MC samples in Sections 7.1 and 7.2. Event selection criteria are discussed in Section 7.3. Section 7.4 compares basic properties of electrons and jets between data and MC.

7.1 Data Sample

The data sample used for this analysis was collected between April 2002 and June 2004 and contains approximately 876 million events. The raw data are processed with the p14 version of the DØ reconstruction software. In order to create a final data sample of manageable size, additional pre-selection or *skimming* criteria are applied:

EM1TRK skimming Each event in the data set is required to have at least one EM object with $ID = 10$ or ± 11 , $p_T > 8$ GeV, and a track with $p_T > 5$ GeV within $\Delta\phi = 0.1$ of the EM object. These requirements reduce the size of the data sample to approximately 57 million events.

Root-tuple creation The reconstructed data are reformatted into an object oriented ntuple format (*root-tuple*) using the ATHENA (61) software package (version p16-br-03). At this stage, JES corrections (version 5.3) and a calorimeter noise suppression algorithm (T42 (62)) are applied.

Root-tuple skimming The root-tuple data are further skimmed by requiring at least one EM object with EM fraction > 0.9 , Isolation < 0.15 , H-Matrix(7) < 12.0 , $|\eta_{det}| < 1.1$ and a track match¹ in each event. The final analysis root-tuple contains 2.4 million events.

Data flagged as unusable by data-quality experts are excluded from the analysis. SMT, CFT, calorimeter, and luminosity subsystems of the detector are required to be fully operational. Additionally, all data taking periods with limited L1CAL trigger coverage ($|\eta| < 0.8$) are excluded (63).

Events for the analysis are selected based on the requirement that the trigger system identified at least one EM object (*single electron triggers*). Only single electron triggers that were run *unprescaled* are used. When the trigger conditions implemented in the trigger list were not providing the required reduction in event rate, *prescaling factors* were applied. For example, if a prescaling factor of 3 is applied to a given L1 trigger, then only one out of three events satisfying that L1 condition is kept for subsequent trigger evaluation.

The data taking period for this analysis can be divided into two periods during which different lists of single EM triggers were implemented. The following is the prioritized order of trigger combinations for trigger lists before *global_CMT-12* (runs ≤ 178732 , “pre-v12 dataset”)² (64):

- EM_HI_SH or EM_HI_2EM5_SH

¹ χ^2 probability for best track using the distance in η/Φ and E/p

²The statement “Trigger A or Trigger B” refers to the fact that a given event is accepted, if Trigger A and Trigger B are unprescaled, and the trigger requirements for either Trigger A or Trigger B are met.

- EM_HL_SH
- EM_HI
- EM_MX_SH
- EM_MX

The trigger combinations for trigger list *global_CMT-12* (runs ≥ 178722 , “v12 dataset”) are:

- E1_SHT20 or E2_SHT20 or E3_SHT20 or E1_SH30
- E1_SHT20 or E2_SHT20 or E1_SH30
- E1_SHT20 or E1_SH30
- E1_SHT20

Table I contains details of the individual triggers.

A total integrated luminosity of 343 pb^{-1} was available for this analysis after trigger selection and exclusion of unusable data due to bad quality.

7.2 Monte Carlo Samples

7.2.1 PYTHIA and ALPGEN Samples

The MC samples used for data comparisons and acceptance estimations are summarized in Table II. For studies regarding the inclusive $Z/\gamma^* \rightarrow e^+e^-$ cross section, a PYTHIA $Z/\gamma^* \rightarrow e^+e^-$ inclusive sample is used. For higher jet multiplicities, events are generated with ALPGEN and then passed through PYTHIA for parton showering and hadronization.

TABLE I

SINGLE EM TRIGGERS USED IN THIS ANALYSIS			
Trigger	L1	L2	L3
EM_HI_SH	CEM(1,10)	EM(1,12)	ELE_LOOSE_SH_T(1,20)
EM_HI_2EM5_SH	CEM(2,5)	EM(1,12)	ELE_LOOSE_SH_T(1,20)
EM_HI	CEM(1,10)	EM(1,12)	ELE_LOOSE(1,30)
EM_MX_SH	CEM(1,15)	none	ELE_LOOSE_SH_T(1,20)
EM_MX	CEM(1,15)	none	ELE_LOOSE(1,30)
E1_SHT20	CEM(1,11)	none	ELE_NLV_SHT(1,20)
E2_SHT20	CEM(2,6)	none	ELE_NLV_SHT(1,20)
E3_SHT20	CEM(1,9)CEM(2,3)	none	ELE_NLV_SHT(1,20)
E1_SH30	CEM(1,11)	none	ELE_NLV_SH(1,30)

L1 Triggers	
CEM(1,10)	one EM trigger tower with $E_T > 10$ GeV
CEM(2,5)	two EM trigger towers with $E_T > 5$ GeV
CEM(1,15)	one EM trigger tower with $E_T > 15$ GeV
CEM(1,11)	one EM trigger tower with $E_T > 11$ GeV
CEM(2,6)	two EM trigger towers with $E_T > 6$ GeV
CEM(1,9)CEM(2,3)	one EM trigger tower with $E_T > 9$ GeV, another EM trigger tower with $E_T > 3$ GeV

L2 Triggers	
EM(1,12)	one EM candidate with $E_T > 12$ GeV (not present for runs below 169524)

L3 Triggers	
ELE_LOOSE_SH_T(1,20)	one electron with $ \eta < 3.0$ and $E_T > 20$ GeV passing loose requirements including shower shape cuts
ELE_LOOSE(1,30)	one electron with $ \eta < 3.0$ and $E_T > 30$ GeV passing loose requirements
ELE_NLV_SHT(1,20)	one electron with $ \eta < 3.6$ and $E_T > 20$ GeV passing tight shower shape cuts
ELE_NLV_SH(1,30)	one electron with $ \eta < 3.6$ and $E_T > 30$ GeV passing loose shower shape cuts

The electron energy resolution measured in data is not correctly described by the MC simulation. Additional energy smearing is applied to the MC electrons to account for the difference. p_x , p_y , p_z and energy of the electrons are multiplied by $c \cdot \text{Gauss}(1, f)$, where $\text{Gauss}(1, f)$ is the additional smearing parameter which is chosen from a Gaussian distribution with mean 1 and width f , and c is an overall calibration factor. The following values for the smearing parameters are used (65):

- $f = 0.045$
- $c = 1.003$

We also adjust the jet energy resolution in MC to match the jet resolution in data (66). The parameterization of the jet energy resolution is given by:

$$\frac{\sigma(p_T)}{p_T} = \sqrt{\frac{N^2}{p_T^2} + \frac{S^2}{p_T} + C^2}, \quad (7.1)$$

where the constants C , S , and N represent the gain fluctuations, sampling fluctuations, and noise contributions respectively. Table III summarizes all coefficients for different detector regions.

Using the p_T and η_{det} of the MC jets, the data and MC resolutions are calculated. If the data resolution is better than the MC resolution for a given jet, no additional smearing is applied. If the jet resolution in MC is worse than in data, the MC jet energy resolution is adjusted by

applying a multiplicative smearing factor (Equation 7.2) to the 4-vector components of each jet.

$$\text{Smearing Factor} = \text{Gauss} \left(1, \sqrt{\left(\frac{\sigma(p_T)}{p_T} \right)_{data}^2 - \left(\frac{\sigma(p_T)}{p_T} \right)_{MC}^2} \right) \quad (7.2)$$

Process	Generators	Size
$Z(\gamma^*) \rightarrow e^+e^-$	PYTHIA	400k
$Zj \rightarrow e^+e^-j$	ALPGEN + PYTHIA	150k
$Zjj \rightarrow e^+e^-jj$	ALPGEN + PYTHIA	180k
$Zjjj \rightarrow e^+e^-jjj$	ALPGEN + PYTHIA	15k

TABLE II

LIST OF MONTE CARLO SAMPLES

7.2.2 CKKW Samples

Table IV summarizes the CKKW samples used in the analysis. The samples use a matching threshold of $p_T > 15$ GeV. Partons are generated with $|\eta| < 2.5$. The Z boson has a generated mass between 75 GeV and 105 GeV. The matrix element generation with MADGRAPH was done up to jet multiplicities of 3. Higher jet multiplicities are due to parton showering and hadronization simulated with PYTHIA. The factorization scale is set to $\mu_F^2 = M_Z^2$. The renor-

Coefficient	$ \eta_{det} < 0.5$	$0.5 < \eta_{det} < 1.0$	$1.0 < \eta_{det} < 1.5$	$ \eta_{det} > 1.5$
N_{data}	5.05	$9.06 \cdot 10^{-9}$	2.24	6.42
S_{data}	0.753	1.2	0.924	$4.5 \cdot 10^{-10}$
C_{data}	0.0893	0.087	0.135	0.0974
N_{MC}	4.26	4.61	3.08	4.83
S_{MC}	0.658	0.621	0.816	$5.13 \cdot 10^{-7}$
C_{MC}	0.0436	0.0578	0.0729	0.0735

TABLE III

JET ENERGY RESOLUTION PARAMETERS IN DATA AND MC.

malization scale is set to $\mu_R^2 = p_{Tjet}^2$ for jets from initial state radiation and $\mu_R^2 = k_{Tjet}^2$ for jets from final state radiation.

Process	Generators	Size
$Zj \rightarrow e^+e^-j$	MADGRAPH + PYTHIA	234k
$Zjj \rightarrow e^+e^-jj$	MADGRAPH + PYTHIA	20k
$Zjjj \rightarrow e^+e^-jjj$	MADGRAPH + PYTHIA	3k

TABLE IV

LIST OF CKKW SAMPLES

7.2.3 MCFM Cross Sections

MCFM inclusive cross sections are calculated to NLO for up to 2 partons. The kinematic and geometric jet cuts are the same as used in the analysis: jet $p_T > 20$ GeV, $|\eta| < 2.5$. The Z boson has a mass between 80 GeV and 100 GeV, and CTEQ6M was selected for the PDF. The renormalization and factorization scales are set to $\mu_{F/R}^2 = M_Z^2 + p_{TZ}^2$.

7.3 Event Selection

The following selection criteria are applied in order to assure that events with two high p_T electrons contained within the central calorimeter and originating from the decay of a Z/γ^* gauge boson are selected. After identifying these Z *candidate* events, the presence of n high p_T jets is required.

7.3.1 Primary Vertex

The efficiency to reconstruct the PV is about 100% in the central region of the detector, and decreases outside of the SMT fiducial volume. Therefore, the PV is required to be within 60 cm of the detector center along the beam pipe (z-axis).

7.3.2 Electron Selection

EM objects have to satisfy the following requirements:

- Loose electrons:
 - ID = 10 or ± 11
 - EM Fraction > 0.9
 - Isolation < 0.15

- $\text{H-Matrix}(7) < 12$
- $p_T > 25 \text{ GeV}$
- $|\eta_{det}| < 1.1$
- Tight electrons:
 - Requirements of loose electron.
 - Track match¹ with $P(\chi^2) > 0.01$

7.3.3 Z Selection

Z candidates are selected based on the following criteria:

- Two loose electrons.
- At least one of the two electrons needs to be tight.
- One of the two electrons must have fired the trigger.
- Diem invariant mass window cut: $75 \text{ GeV} < M_{ee} < 105 \text{ GeV}$

7.3.4 Jet Selection

Jets are formed using the Run II Midpoint Jet Cone Algorithm with a cone size of 0.5 and are selected based on the following criteria:

- $0.05 < \text{EMF} < 0.95$
- $\text{HotF} < 10$.

¹ χ^2 probability for best track using the distance in η/Φ and E/p

- $N_{90} > 1$.
- $CHF < 0.4$
- L1 confirmation
- JES corrected $p_T > 20$ GeV
- $|\eta_{physics}| < 2.5$
- Since the jet algorithm identifies *fake* jets originating from electron energy deposits, all jets overlapping with electrons coming from the Z/γ^* boson within $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$ of 0.4 are removed.

7.3.5 Event Statistics

Table V summarizes the number of $Z/\gamma^* \rightarrow e^+e^-$ event candidates for different exclusive jet multiplicities.

7.4 Data vs Monte Carlo

This section presents a comparison of basic kinematic distributions for electrons, Z candidates, and jets between data and MC simulations¹. It is important that the MC distributions describe the data distributions as accurately as possible. The MC simulations are used to account for the fraction of events that are lost due to kinematic and geometric electron cuts, the diem invariant mass cut, and the primary vertex cut (*acceptance*).

¹The MC distributions are normalized to the number of events in data.

Sample	N	Fraction
$Z/\gamma^* + 0$ jets	12,247	0.8815
$Z/\gamma^* + 1$ jets	1,427	0.1027
$Z/\gamma^* + 2$ jets	189	0.0136
$Z/\gamma^* + 3$ jets	25	0.0018
$Z/\gamma^* + 4$ jets	3	0.0002
$Z/\gamma^* + 5$ jets	2	0.0001
Total	13,893	1.0000

TABLE V

EVENT BREAKDOWN BY EXCLUSIVE JET MULTIPLICITIES ASSOCIATED WITH Z/γ^* PRODUCTION BEFORE ANY BACKGROUND IS SUBTRACTED OR ANY CORRECTIONS ARE APPLIED.

7.4.1 Primary Vertex Comparison

Figure 32 compares the primary vertex distribution between data and inclusive $Z/\gamma^* \rightarrow e^+e^-$ PYTHIA MC.

7.4.2 Z p_T Comparisons

Figure 33 shows the Z p_T comparison between data and PYTHIA MC. Since PYTHIA is a LO ($2 \rightarrow 2$) generator at the hard process, there is disagreement in the Z p_T distribution (especially at high p_T) between data and MC. To account for this discrepancy, an additional correction based on the Z p_T comparison between data and MC is applied. The Z p_T correction is also shown in Figure 33.

Figure 34 and Figure 35 show Z p_T comparisons when using ALPGEN + PYTHIA for $Z+1$ jet and $Z+2$ jet samples. The agreement between ALPGEN + PYTHIA MC and data




Figure 32. Primary vertex distribution in data and MC (PYTHIA) for the inclusive sample.
The MC distribution is normalized to the number of events in data.




Figure 33. Comparison of $Z p_T$ between data and PYTHIA MC (left), and ratio correction factor (right) for the inclusive sample. The MC distribution (left) is normalized to the number of events in data.

is deemed acceptable. Therefore, no additional $Z p_T$ correction is applied to the ALPGEN + PYTHIA MC samples.




Figure 34. Comparison of Z p_T between data and ALPGEN + PYTHIA $Z+1$ jet MC. The MC distribution is normalized to the number of events in data.




Figure 35. Comparison of Z p_T between data and ALPGEN + PYTHIA $Z+2$ jets MC. The MC distribution is normalized to the number of events in data.

7.4.3 $Z/\gamma^* (\rightarrow e^+e^-) + \geq n$ Jet Comparisons

7.4.3.1 $Z/\gamma^* (\rightarrow e^+e^-)$ Inclusive Sample

This section compares basic kinematic distributions for electrons and Z candidates after applying all corrections (Trigger, EM, Tracking, Z p_T - see Chapters 8.1 and 9.1 for a description of the corrections). Figure 36 compares basic electron and Z kinematic distributions. Figure 37 compares the diem invariant mass distribution. The average Z mass is 91.02 GeV with a width of 4.03 GeV.

7.4.3.2 $Z/\gamma^* (\rightarrow e^+e^-) + \geq 1$ Jet Sample

Figure 38 shows comparisons of basic electron and Z distributions. Figure 39 shows a comparison of the diem invariant mass peak. Figure 40 and Figure 41 show comparisons of basic kinematic distributions for the leading p_T jet.

7.4.3.3 $Z/\gamma^* (\rightarrow e^+e^-) + \geq 2$ Jet Sample

Figure 42 shows comparisons of basic electron and Z distributions. Figure 43 shows a comparison of the diem invariant mass peak and Figure 44, Figure 45 and Figure 46 show comparisons of basic kinematic distributions for the two laeding p_T jets.




Figure 36. p_T of both Z electrons (top left), physics η of both Z electrons (bottom left), Z p_T (top right), Z rapidity (bottom right) for the $Z/\gamma^* \rightarrow e^+e^-$ inclusive sample in data and MC (PYTHIA). The MC distribution is normalized to the number of events in data.




Figure 37. Dielectron invariant mass comparison for the $Z/\gamma^* \rightarrow e^+e^-$ inclusive sample in data and MC (PYTHIA). Data is background subtracted. The MC distribution is normalized to the number of events in data.




Figure 38. p_T of both Z electrons (top left), physics η of both Z electrons (bottom left), Z p_T (top right), Z rapidity (bottom right) for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1 \text{ jet}$ sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.




Figure 39. Dielectron invariant mass comparison for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample in data and MC (ALPGEN). Data is background subtracted. The MC distribution is normalized to the number of events in data.




Figure 40. p_T (linear and logarithmic), physics η and physics Φ of all jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1 \text{ jet}$ sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.




Figure 41. p_T (linear and logarithmic), physics η and physics Φ of the leading jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1 \text{ jet}$ sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.




Figure 42. p_T of both Z electrons (top left), physics η of both Z electrons (bottom left), Z p_T (top right), Z rapidity (bottom right) for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2 \text{ jet}$ sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.




Figure 43. Dielectron invariant mass comparison for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). Data is background subtracted. The MC distribution is normalized to the number of events in data.




Figure 44. p_T (linear and logarithmic), physics η and physics Φ of all jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2 \text{ jet}$ sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.




Figure 45. p_T (linear and logarithmic), physics η and physics Φ of the leading jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2 \text{ jet}$ sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.




Figure 46. p_T (linear and logarithmic), physics η and physics Φ of the next to leading jets for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample in data and MC (ALPGEN). The MC distributions are normalized to the number of events in data.

CHAPTER 8

MEASUREMENT OF THE $Z/\gamma^*(\rightarrow e^+e^-)$ INCLUSIVE CROSS SECTION

In order to determine the $Z/\gamma^*(\rightarrow e^+e^-)$ inclusive cross section, the following equation is evaluated:

$$\sigma \times \text{BR}(Z/\gamma^* \rightarrow e^+e^-) = \frac{N - B}{\mathcal{L} \times \varepsilon}. \quad (8.1)$$

$N - B$ is the number of signal events which is estimated from the diem invariant mass histogram after background subtraction. The primary background is from multi-jet production from QCD processes in which the jets have a large electromagnetic component or are mismeasured in such a way that the jets pass the electron selection criteria. The shape of the QCD background in the diem invariant mass histogram follows an exponential distribution.

The goal is to measure the cross section for diem pairs where both γ^* (*Drell-Yan*) and Z boson exchange contribute. Contributions from pure Z boson decays will show up as a peak around the Z mass at ≈ 91 GeV in the diem invariant mass histogram. The Drell-Yan component follows an exponential distributions.

The total integrated luminosity of the dataset \mathcal{L} is approximately 343 pb^{-1} .

The efficiency ε to measure e^+e^- pairs resulting from Z/γ^* decays can be factorized according to:

$$\varepsilon_{tot} = \varepsilon_{trigger} \cdot \varepsilon_{EM} \cdot \varepsilon_{track} \cdot \varepsilon_{acceptance} \quad (8.2)$$

where $\varepsilon_{trigger}$ is the efficiency of the event to have at least one electron to pass all trigger levels, ε_{EM} is the efficiency of reconstructing two EM clusters which pass all electron ID cuts, ε_{track} is the efficiency of finding at least one track which matches an EM cluster and $\varepsilon_{acceptance}$ is the efficiency of the kinematic and geometric electron cuts, the diem invariant mass cut and the primary vertex cut. Practically, all efficiencies are applied (except for $\varepsilon_{acceptance}$) as corrections to the diem invariant mass histogram. This procedure is outlined in the following chapters.

8.1 Efficiencies

8.1.1 Trigger Efficiency

The combined efficiency of all triggers per electron is studied with a *tag-and-probe method* using Z candidate events with invariant mass between 70 and 110 GeV. For this method, both Z candidate electrons are considered as possible “tags”. An electron becomes a “tag” if it passes trigger requirements for at least one unscaled trigger in the trigger combination. To pass the requirements of a trigger, an electron must have a matching trigger object at each level which passes all cuts for the corresponding trigger. Both the tag and probe electrons must satisfy the following requirements:

- $p_T > 20$ GeV
- EM Fraction > 0.9
- Isolation < 0.15
- H-Matrix(7) < 12
- Track match with $P(\chi^2) > 0.01$

The probe electron must have matching trigger objects at L1, L2 and L3 within $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$ of 0.4.

Trigger efficiencies are parameterized versus EM object p_T and derived separately for pre-v12 and v12 data. In cases where the L2 subsystem was not operative (all runs with run numbers $\leq 169,523$) only L1 and L3 trigger objects were used. The average trigger efficiencies for the pre-v12 and v12 datasets are (with statistical errors):

- $\varepsilon_{pre-v12}(\text{Trigger}) = 94.6\% \pm 0.3\%$
- $\varepsilon_{v12}(\text{Trigger}) = 98.2\% \pm 0.1\%$

Figure 47 shows the parameterized trigger efficiencies for both datasets.

Trigger inefficiencies are corrected in the following way:

- In a given event the trigger efficiency curves are used to get the trigger efficiencies ε_1 and ε_2 for the two EM objects (based on p_T).
- To get the event based trigger efficiency, all permutations for the two EM objects to fire a trigger need to be taken into account:

$$\varepsilon_{trigger} = \varepsilon_1 \cdot (1 - \varepsilon_2) + \varepsilon_2 \cdot (1 - \varepsilon_1) + \varepsilon_1 \cdot \varepsilon_2 = \varepsilon_1 + \varepsilon_2 - \varepsilon_1 \cdot \varepsilon_2 \quad (8.3)$$

- The inverse of $\varepsilon_{trigger}$ is applied as a corrective weight when the diem invariant mass histogram is filled.
- The corrected diem invariant mass histogram is used to derive the number of signal events for the cross section calculation.




Figure 47. Trigger efficiencies for pre-v12 (top) and v12 (bottom) datasets vs EM object p_T

8.1.2 EM Reconstruction and Identification Efficiency

To determine EM efficiencies, a tag and probe method is used. The tag leg consists of an electron candidate, and the probe leg consists of a track. The tag electron must pass all the loose electron selection cuts, have a matched track and satisfy trigger requirements for the event. Both tag and probe tracks must satisfy the following selection criteria (67):

- Stereo track¹
- $25 \text{ GeV} < p_T < 80 \text{ GeV}$
- χ^2 probability for best track < 8 (using the distance in η/Φ and E/p)
- Distance of closest approach between track and beam position in the R - Φ plane $< 0.3 \text{ cm}$
- Δz_{vertex} of the two tracks $< 4 \text{ cm}$
- $|\eta_{\text{detector}}| < 1.1$

Tag electron selection criteria:

- $ID = 10 \text{ or } \pm 11$
- $EMFraction > 0.9$
- $Isolation < 0.15$
- $H\text{-Matrix}(7) < 12$
- $p_T > 25 \text{ GeV}$
- $|\eta_{\text{det}}| < 1.1$
- No fiducial restrictions in ϕ .
- Matched with tag track within $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = \sqrt{0.1^2 + 0.1^2} = 0.14$.
- Must have fired the trigger.

¹Requiring hits in stereo layers of the tracking system (see Chapter 3.3.2).

Possible background contamination is reduced by requiring that tag- and probe-tracks have opposite signs, and by imposing a cut on the missing transverse energy of the event (missing $E_T < 15$ GeV). The following lists additional requirements:

- $|PVZ| < 60$ cm
- Tag-electron-probe-track invariant mass cut: $70 \text{ GeV} < M_{ee} < 110 \text{ GeV}$

Once an event is found which satisfies all of the above requirements, a denominator histogram is filled. Probe track ϕ and p_T histograms are used for parameterized efficiencies. Tag-electron-probe-track diem invariant mass histograms are used for average efficiencies. If a reconstructed EM cluster is found nearby the probe-track ($\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = \sqrt{0.1^2 + 0.1^2} = 0.14$) which passes the EMID cuts (HMx, EMF, Iso), the respective numerator histogram is filled.

Since average efficiencies are derived from diem invariant mass histograms, an additional step to subtract background can be applied. The background in the sidebands of the diem invariant mass distributions is estimated, and extrapolated into the signal region where it is subtracted. The average EM reco and ID efficiencies in data and MC are (with statistical errors):

- $\varepsilon_{data}(\text{EM}) = 88.9\% \pm 0.3\%$
- $\varepsilon_{MC}(\text{EM}) = 93.1\% \pm 0.1\%$

The sideband background subtraction cannot be applied in the case of parameterized efficiencies. The level of background contamination is examined by deriving the average efficiency in data without the sideband background subtraction. The result is within 1% of the sideband

subtracted value: $88.2\% \pm 0.2\%$.

Figure 48 and Figure 49 show the EM efficiencies for data and MC in a one-dimensional parameterization versus probe track Φ and p_T . Note that the central calorimeter has narrow uninstrumented regions between the azimuthal module boundaries (*phi cracks*). An EM object entering the calorimeter near these boundaries can lose a portion of its energy in these cracks, which results in decreased EM efficiencies for these regions.

EM inefficiencies are corrected in the following way:

- In a given event two-dimensional efficiency curves are used (versus p_T and Φ) to estimate the EM efficiencies ε_1 and ε_2 for the two EM objects (based on their p_T and Φ).
- To estimate the event based EM efficiency, the product of ε_1 and ε_2 is taken:

$$\varepsilon_{EM} = \varepsilon_1 \cdot \varepsilon_2 \tag{8.4}$$

- The inverse of ε_{EM} is applied as a corrective weight when filling the diem invariant mass histogram.

8.1.3 EM-Track Match Efficiency

Average track finding and matching efficiencies are derived using diem invariant mass histograms (Figure 50 to Figure 53).

Using a convolution of a Gaussian and Breit-Wigner fit for the Z peak and an exponential shape to describe the QCD and Drell-Yan contributions, the number of events under the Z peak is extracted from the four diem invariant mass histograms: $N_{1trk}(data)$, $N_{2trk}(data)$,




Figure 48. EM efficiencies versus probe track Φ and p_T in data. The Φ distribution shows the $\text{modulus}(\Phi, \frac{2\pi}{32})$ distribution to illustrate the effect of the calorimeter Φ -module boundaries.




Figure 49. EM efficiencies versus probe track Φ and p_T in MC. The Φ distribution shows the modulus($\Phi, \frac{2\pi}{32}$) distribution to illustrate the effect of the calorimeter Φ -module boundaries.

$N_{1trk}(MC)$ and $N_{2trk}(MC)$. $N_{1trk}(data)$ and $N_{1trk}(MC)$ are the number of Z candidates when requiring at least one track match in data and MC; $N_{2trk}(data)$ and $N_{2trk}(MC)$ are the number of Z candidates when requiring exactly two track matches in data and MC. These numbers are used to estimate the average track finding and matching efficiencies in data and MC:

$$\varepsilon_{data}(\text{Tracking}) = \frac{2 \cdot N_{2trk}(data)}{N_{2trk}(data) + N_{1trk}(data)} = 77.1\% \pm 0.3\% \quad (8.5)$$

$$\varepsilon_{MC}(\text{Tracking}) = \frac{2 \cdot N_{2trk}(MC)}{N_{2trk}(MC) + N_{1trk}(MC)} = 87.8\% \pm 0.03\% \quad (8.6)$$

The tracking inefficiencies are corrected in the following way:

- In each event the average tracking efficiency $\varepsilon_{average}$ is used (Equation 8.5 and Equation 8.6).
- To get the event based tracking efficiency, all permutations for one or two tracks need to be taken into account:

$$\varepsilon_{tracking} = 2 \cdot \varepsilon_{average}(1 - \varepsilon_{average}) + \varepsilon_{average}^2 = 2 \cdot \varepsilon_{average} - \varepsilon_{average}^2 \quad (8.7)$$

- The inverse of $\varepsilon_{tracking}$ is applied as a corrective weight when filling the diem invariant mass histogram.

8.1.4 Acceptance

The $Z/\gamma^* \rightarrow e^+e^-$ PYTHIA MC sample is used to estimate the acceptance. The numerator contains the number of events satisfying the following requirements at the reconstructed level:




Figure 50. Invariant mass in data when requiring at least one track-matched electron.




Figure 51. Invariant mass in data when requiring two track-matched electrons.




Figure 52. Invariant mass in MC when requiring at least one track-matched electron.




Figure 53. Invariant mass in MC when requiring two track-matched electrons.

- Primary vertex cut: $|PVZ| < 60$ cm
- Electron cuts: $p_T > 25$ GeV and $|\eta| < 1.1$
- Diem invariant mass cut: $75 \text{ GeV} < M_{ee} < 105 \text{ GeV}$

The denominator contains the number of events with generated Z/γ^* particles that are within the diem invariant mass window.

The acceptance with statistical error for inclusive $Z/\gamma^* \rightarrow e^+e^-$ is:

$$\text{Acceptance}(Z/\gamma^* \rightarrow e^+e^- + X) = 21.4\% \pm 0.1\% \quad (8.8)$$

Since the numerator in the acceptance calculation involves two reconstructed electrons, a corrective weight is applied to the numerator. Based on the p_T and Φ values of the two electrons, their reconstruction efficiencies are estimated to be $\approx 98\%$. The product of the inverse of those reconstruction efficiencies yields the corrective weight. The Z p_T correction factor is also applied (see Chapter 7.4.2) as an additional weight in both numerator and denominator of the acceptance.

8.2 Cross Section Calculation

After applying all corrections, the number of corrected signal events is estimated from the diem invariant mass distribution (Figure 54). A convolution of a Gaussian and Breit-Wigner shape is fitted to the Z peak. An exponential shape is used to describe the QCD and Drell-Yan contributions. The choice of using an exponential shape for the QCD background is justified

by examining the diem invariant mass distribution of EM object pairs that were selected by applying “anti-electron cuts”:

- All criteria that are applied to loose electron candidates as described in Section 7.3.2 except for the H-Matrix cut.
- $\text{H-Matrix}(7) > 35$
- Two of these objects per event.

In this way, two jets are selected with high electromagnetic energy content in the shower.

Since the Drell-Yan component is part of the signal, the QCD component needs to be disentangled from the Drell-Yan component. Using the inclusive $Z/\gamma^* \rightarrow e^+e^-$ PYTHIA MC sample, the percentage of Drell-Yan events in $Z/\gamma^* \rightarrow e^+e^-$ decays is estimated by fitting a Gaussian and Breit-Wigner shape to the Z component and an exponential shape to the Drell-Yan component. 2.06% of the events in the inclusive $Z/\gamma^* \rightarrow e^+e^-$ sample are due to Drell-Yan.

Based on these fits the number of signal events from direct Z Boson and Drell-Yan decays is extracted, as well as the number of QCD background events in the diem invariant mass signal window ($75 \text{ GeV} < M_{ee} < 105 \text{ GeV}$) ¹:

- Number of corrected signal events from Z Boson and Drell-Yan decays = 18,263.8
- Number of corrected QCD background events = 407.5

¹The number of signal events is derived by counting all entries for a particular M_{ee} bin and subtracting from it the number of entries from the background fit.

Based on the integrated luminosity (343 pb^{-1}) and the acceptance (21.4%), the inclusive Z/γ^* production cross section times branching fraction into electrons is calculated:

$$\sigma \times \text{BR}(Z/\gamma^* \rightarrow e^+e^-) = \frac{N - B}{\mathcal{L} \times A} = 248.9 \pm 2.5(\text{stat}) \text{ pb} \quad (8.9)$$

In order to check the procedure that leads to the cross section measurement, a MC closure test is performed. The number of signal events, acceptance, and luminosity are evaluated in MC. This leads to a calculated MC cross section (179 pb) that is being compared to the input cross section that was used to generate the MC sample (183 pb). The calculated MC cross section is in good agreement with the MC input cross section.

8.3 Comparisons to Other Measurements

Using the inclusive PYTHIA MC sample, correction factors are derived which allow to estimate the Z/γ^* inclusive cross section for different diem invariant mass windows ¹. To calculate the correction factors, the number of Z/γ^* candidates is counted at the particle level within different diem invariant mass ranges. The ratio is taken which yields the correction that is needed to account for the change in the diem invariant mass range.

Using the cross section ratio between the inclusive Z/γ^* cross section and the inclusive Z cross section in MC, another correction factor is derived which allows to convert the measured inclusive Z/γ^* cross section into the inclusive Z cross section (68).

¹The MC sample used a generator cut of $|\eta_Z| < 4.2$. Although no restriction in η_Z would have been preferable, the impact on the final results is believed to be negligible.




Figure 54. Diem invariant mass distribution for $Z/\gamma^* \rightarrow e^+e^- + X$ (Mean = $91.02 \text{ GeV} \pm 0.04 \text{ GeV}$, Width $4.03 \text{ GeV} \pm 0.04 \text{ GeV}$).

Process	Mass Range	Conversion
(a) Z/γ^*	$75 \text{ GeV} < M_{ee} < 105 \text{ GeV}$	248.9 pb
(b) Z/γ^*	$66 \text{ GeV} < M_{ee} < 116 \text{ GeV}$	$248.9 \text{ pb} \cdot \frac{392,174}{379,255} = 257.4 \text{ pb}$
(c) Z/γ^*	$60 \text{ GeV} < M_{ee} < 130 \text{ GeV}$	$248.9 \text{ pb} \cdot \frac{400,000}{379,255} = 262.5 \text{ pb}$
(d) Z	$60 \text{ GeV} < M_{ee} < 130 \text{ GeV}$	$248.9 \text{ pb} \cdot \frac{1.852}{1.869-0.948} = 260.2 \text{ pb}$

TABLE VI

CROSS SECTION RESULTS FOR DIFFERENT DIEM INVARIANT MASS RANGES AND PROCESSES.

Table VI lists the inclusive Z/γ^* cross sections for different mass ranges, as well as the conversion into the inclusive cross section for pure Z boson exchange. Result (b) is compared with the CDF measurement (69) for the inclusive Z/γ^* cross section ($66 \text{ GeV} < M_{ee} < 116 \text{ GeV}$) of $255.8 \pm 3.9(\text{stat}) \text{ pb}$.

CHAPTER 9

MEASUREMENT OF THE $Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ JET CROSS SECTIONS

This section outlines the procedure to measure the $Z/\gamma^*(\rightarrow e^+e^-)$ production cross section for different inclusive jet multiplicities. For each jet multiplicity the number of signal events is determined from the diem invariant mass histograms. All efficiencies are examined for jet multiplicity dependence and applied to the diem invariant mass distributions as corrections (acceptance corrections are applied separately). Jet reconstruction inefficiencies are taken into account, as well as jet acceptance losses due to the electron-jet-overlap cut. The final cross sections are quoted for particle level jets (as opposed to calorimeter jets). The *unsmearing* procedure is described that is used to convert calorimeter level jet multiplicities to particle level jet multiplicities. This is done in order to obtain final cross sections that are independent from detector resolution effects.

9.1 Efficiencies vs Jet Multiplicity

In the following sections the PYTHIA MC sample is used to derive corrections for the inclusive sample, while ALPGEN MC samples are used for the n-jet corrections.

9.1.1 Trigger Efficiency

No significant variations in the trigger efficiencies are observed as jet activity increases (Figure 55). Therefore the same trigger corrections as for the inclusive sample are applied (see Chapter 8.1.1). Table VII summarizes electron trigger efficiencies for the pre-v12 and




Figure 55. Average object based trigger efficiencies in data versus inclusive jet multiplicity.

v12 datasets for different inclusive jet multiplicities. An additional systematic error of $\pm 5\%$ is assigned for all jet multiplicities.

9.1.1.2 EM Reconstruction and Identification Efficiency

The same EM corrections as for the inclusive sample are applied to each jet multiplicity sample (see Chapter 8.1.2). Residual inefficiencies due to additional jet activity are estimated by examining the distribution of average electron EM reco and ID efficiencies versus jet multiplicity

Jet multiplicity	pre-v12	v12
≥ 0	$94.6\% \pm 0.3\% \pm 5.0\%$	$98.2\% \pm 0.1\% \pm 5.0\%$
≥ 1	$93.1\% \pm 1.0\% \pm 5.0\%$	$96.9\% \pm 0.5\% \pm 5.0\%$
≥ 2	$95.2\% \pm 2.1\% \pm 5.0\%$	$95.5\% \pm 1.7\% \pm 5.0\%$

TABLE VII

OBJECT BASED TRIGGER EFFICIENCIES WITH STATISTICAL AND SYSTEMATIC ERRORS (RESPECTIVELY) FOR THE PRE-V12 AND V12 DATASETS FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES.

(Figure 56). The averaged efficiencies are derived using the procedure outlined in Chapter 8.1.2 for data and MC samples of different jet multiplicities.

Table VIII summarizes the electron EM reco and ID efficiencies in data and MC for different jet multiplicities. No significant change of the average efficiencies with respect to jet multiplicity is observed in data. Therefore no additional correction is applied. Due to the variation of the various efficiencies a systematic error of $\pm 3\%$ is assigned for all jet multiplicities.

Based on the efficiency drop in MC a factor of 1.02^2 is applied as a corrective weight to each event. This value is derived by taking the ratio of the EM efficiency for the inclusive sample and the average of the EM efficiencies for the 1-jet, 2-jet and 3-jet samples.

9.1.3 EM-Track Match Efficiency

Figure 57 and Table IX show the average object based tracking efficiencies for different jet multiplicities. In MC, the value from the inclusive sample is used to correct for tracking inefficiencies for all jet multiplicities. In data, the inclusive value is used for the inclusive sample,




Figure 56. Average object based EM reco and ID efficiencies in data and MC versus inclusive jet multiplicity. There was not enough statistics available to estimate the EM efficiency in data for ≥ 3 jets.

Jet multiplicity	data	MC
≥ 0	$88.9\% \pm 0.3\% \pm 3.0\%$	$93.1\% \pm 0.1\%$
≥ 1	$87.2\% \pm 1.0\% \pm 3.0\%$	$92.3\% \pm 0.3\%$
≥ 2	$90.0\% \pm 2.5\% \pm 3.0\%$	$91.2\% \pm 1.0\%$
≥ 3	(n/a)	$90.1\% \pm 3.5\%$

TABLE VIII

OBJECT BASED EM RECO AND ID EFFICIENCIES WITH STATISTICAL AND SYSTEMATIC ERRORS (RESPECTIVELY) IN DATA AND MC FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES. THERE WAS NOT ENOUGH STATISTICS AVAILABLE TO ESTIMATE THE EM EFFICIENCY IN DATA FOR ≥ 3 JETS.

Jet multiplicity	data	MC
≥ 0	$77.1\% \pm 0.3\%$	$87.8\% \pm 0.03\%$
≥ 1	$74.5\% \pm 0.9\%$	$87.7\% \pm 0.3\%$
≥ 2	$72.1\% \pm 2.5\%$	$87.5\% \pm 0.9\%$

TABLE IX

OBJECT BASED TRACKING EFFICIENCIES WITH STATISTICAL ERRORS IN DATA AND MC FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES.

the 1-jet value is used for the 1-jet sample and the 2-jet value is used for all other jet samples.

Table X lists the systematic errors for the data efficiencies. For the 1-jet and 2-jet samples, we use the respective statistical errors as systematics. For the 3-jet, 4-jet and 5-jet samples, the 2-jet statistical error is added in quadrature with the difference between a linear fit to the available data points and the 2-jet efficiency value.




Figure 57. Average object based tracking efficiencies in data and MC versus inclusive jet multiplicity.

Jet multiplicity	Data Efficiency	Systematic Uncertainty
≥ 0	77.1%	$\pm 0.3\%$
≥ 1	74.5%	$\pm 0.9\%$
≥ 2	72.1%	$\pm 2.5\%$
≥ 3	72.1%	$\pm 3.5\%$
≥ 4	72.1%	$\pm 5.6\%$
≥ 5	72.1%	$\pm 7.9\%$

TABLE X

OBJECT BASED TRACKING EFFICIENCIES WITH SYSTEMATIC ERRORS.

9.1.4 Acceptance

ALPGEN MC samples are used to estimate the acceptances for different jet multiplicities¹.

The numerator for the n -jet acceptance contains the number of events satisfying the following requirements:

- Primary vertex cut: $|PVZ| < 60$ cm
- Electron cuts: $p_T > 25$ GeV and $|\eta| < 1.1$
- Diem invariant mass cut: $75 \text{ GeV} < M_{ee} < 105 \text{ GeV}$
- Particle level jet cut: n jets with $p_T > 20$ GeV and $|\eta| < 2.5$

The denominator for the n -jet acceptance contains the number of events satisfying the following requirements:

¹The $Z+3$ jet sample is used for jet multiplicities of 3, 4, and 5.

Jet multiplicity	Acceptance
≥ 0	$21.4\% \pm 0.1\%$
≥ 1	$25.1\% \pm 0.2\%$
≥ 2	$25.4\% \pm 0.2\%$
≥ 3	$27.4\% \pm 0.3\%$
≥ 4	$28.5\% \pm 0.7\%$
≥ 5	$30.3\% \pm 1.9\%$

TABLE XI

ACCEPTANCES WITH STATISTICAL ERRORS FOR DIFFERENT JET MULTIPLICITIES.

- MC generator diem invariant mass cut: $75 \text{ GeV} < M_{ee} < 105 \text{ GeV}$
- Particle level jet cut: n jets with $p_T > 20 \text{ GeV}$ and $|\eta| < 2.5$

No additional Z p_T correction is needed since the Z p_T distributions between data and ALPGEN MC agree well (see Chapter 7.4.2). Table XI summarizes the acceptances for different jet multiplicities. On average, higher jet multiplicities lead to higher Z p_T , since the Z boson recoils against the jet(s) in the event. This in turn leads to electrons coming from Z decays that are more likely to pass the acceptance requirements. Therefore, as jet multiplicities increase, acceptances increase as well.

9.1.5 Jet Reconstruction and Identification Efficiency

To estimate the jet reco/ID efficiency in data, the “ Z p_T balance” method is used to measure the scaling factor needed to adjust the MC jet reco/ID efficiency to data. MC is used to measure the “straight” reco/ID jet efficiency with respect to the particle level jets (70). The Z p_T balance

method relies on the recoil of a jet against the Z boson. Events with Z candidates are selected and searched for a recoiling calorimeter jet opposite in Φ . The “efficiency” of finding a recoiling jet is measured as a function of the Z p_T in data and MC. The ratio of these efficiencies in data and MC gives the scaling factor that is applied to the MC in order to make it agree with the data jet reco/ID efficiency. Following this step, the straight jet reco/ID efficiency is measured in MC by matching particle level jets with calorimeter jets within a search cone of $\Delta R = 0.4$. The jet reco/ID efficiency is plotted versus MC particle jet p_T . The p_T values of the particle jets are smeared with the data energy resolutions (see Chapter 7.2.1). Figure 58 shows the data jet reconstruction efficiencies for different regions in the calorimeter.




Figure 58. Jet reco/ID efficiencies in data. CC = $-0.7 < |\eta_{det}| < 0.7$, ICR = $0.7 < |\eta_{det}| < 1.5$, EC = $1.5 < |\eta_{det}| < 2.5$.

9.2 Cross Section Calculation

9.2.1 Unsmearing

In order to determine particle level cross sections, the measured data jet multiplicities are unsmearred from detector resolution effects. A special $Z+1$ jet PYTHIA MC sample is used by choosing PYTHIA 2-to-2 subprocesses 15 and 30¹ with a parton p_T cut-off of 8 GeV and by applying vertex smearing. This sample only contains particle level jets (no detector simulation). The full data resolution smearing is applied to the particle jet p_T as well as the data jet reco/ID efficiencies. Comparing the inclusive jet multiplicity for this MC sample with the one in data gives an idea of how well the MC describes the data (Figure 59). There is increasing disagreement at higher jet multiplicities, since PYTHIA does not include higher order contributions at the hard scatter level. This discrepancy is corrected for by taking the ratio between data and MC for each inclusive jet multiplicity (Figure 60) and then applying correction factors (per event) based on the absolute values of these ratios. After this additional step the inclusive jet multiplicity spectrum in MC is again compared with the one in data with much better agreement (Figure 61). This corrected MC sample is used to derive the coefficients that unsmear the measured data jet multiplicities, also taking into account jet reco/ID inefficiencies. Figure 62 - Figure 73 compare jet p_T and η distributions between the data and MC sample.

To calculate the coefficients, the inclusive jet multiplicity histogram for particle level jets with $p_T > 20$ GeV and $|\eta_{physics}| < 2.5$ is divided by the inclusive jet multiplicity histogram for

¹ $f_i \bar{f}_i \rightarrow gZ^0$ and $f_i g \rightarrow f_i Z^0$




Figure 59. Comparing inclusive jet multiplicities between data and particle level MC (applying data resolution smearing and data jet reco/ID efficiencies). The distributions are normalized with respect to the first bin.

particle level jets with smeared $p_T > 20$ GeV and $|\eta_{physics}| < 2.5$ (after applying jet reco/ID efficiencies). The exact values of these ratios yield the unsmeared and jet reco/ID coefficients which can be applied as multiplicative factors to the measured jet multiplicities in data. Figure 74 shows the numerator and denominator jet multiplicity histograms, as well as the ratio when applying jet smearing and jet reco/ID efficiencies in the denominator. For comparison Figure 75 shows the same distributions without applying jet reco/ID efficiencies, i.e. applying only jet smearing.

The statistical error of each unsmeared and jet reco/ID coefficient is used as a statistical error for the final cross sections (see Chapter 10.7). Table XII summarizes all coefficients.




Figure 60. Ratio of MC (with smearing and jet reco/ID efficiencies) inclusive jet multiplicities and data inclusive jet multiplicities.




Figure 61. Comparing inclusive jet multiplicities between data and particle level MC (after applying correction factors). The distributions are normalized with respect to the first bin.




Figure 62. Comparing jet p_T for all jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 63. Comparing jet p_T for all jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 64. Comparing jet p_T for leading jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.




Figure 65. Comparing jet p_T for leading jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.




Figure 66. Comparing jet p_T for second leading jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.




Figure 67. Comparing jet p_T for second leading jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied). The gray band shows the uncertainty due to the jet energy scale.




Figure 68. Comparing jet p_T for third leading jets between data and particle level MC on a linear scale (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 69. Comparing jet p_T for third leading jets between data and particle level MC on a logarithmic scale (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 70. Comparing jet η for all jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 71. Comparing jet η for leading jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 72. Comparing jet η for second leading jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 73. Comparing jet η for third leading jets between data and particle level MC (with data resolution smearing and jet reco/ID efficiencies applied).




Figure 74. Unsmearing and jet reco/ID particle jet multiplicities (left) and coefficients (right).




Figure 75. Unsmeared jet multiplicities (left) and coefficients (right) without applying jet reco/ID efficiencies.

Jet multiplicity	Unsmearing and jet reco/ID coefficient
≥ 1	1.10 $^{+0.08}_{-0.06}$
≥ 2	1.26 $^{+0.18}_{-0.16}$
≥ 3	1.50 $^{+0.25}_{-0.24}$
≥ 4	1.90 $^{+0.52}_{-0.39}$
≥ 5	4.00 $^{+3.42}_{-1.13}$

TABLE XII

UNSMEARING AND JET RECO/ID COEFFICIENTS WITH SYSTEMATIC UNCERTAINTY DUE TO RESOLUTION AND JET RECO/ID EFFICIENCY.

9.2.2 Electron-Jet-Overlap Correction

Using the tag-and-probe method outlined in Chapter 8.1.2, the ΔR distribution between probe tracks and reconstructed jets that pass all jet quality cuts except for the electron-jet-overlap cut is plotted in data and MC (Figure 76 and Figure 77).

There is an excess of entries at ΔR values of 0 and π due to fake jets (i.e. originated from the electron energy deposits) which survived the jet quality cuts. Therefore, all jets are rejected that are near either of the two electrons within $\Delta R=0.4$. Figure 78 shows the same distribution as in Figure 76 after adding the electron-jet-overlap cut. For comparison Figure 79 shows the ΔR between generated electrons and partons in data and MC.

An additional correction needs to be derived in order to account for the real jets that are removed by the electron-jet-overlap cut. Using the same MC sample as for the unsmearing studies (see Chapter 9.2.1), the correction factors due to the electron-jet-overlap are estimated by taking the ratio of the inclusive parton multiplicity distribution for all partons with $p_T > 20$ GeV and $|\eta| < 2.5$ and the inclusive parton multiplicity distribution for partons that are outside of the ΔR cone with respect to the electrons from the Z .

Correction factors are derived using ΔR cones of size 0.4 and 0.7 and take the middle value as the final correction factors. This is done in order to account for the position resolution between partons and calorimeter jets (see Figure 80). Additionally, a linear fit is applied to the final values in order to compensate for statistical fluctuations.

Table XIII summarizes the electron-jet-overlap correction factors for different jet multi-




Figure 76. ΔR between probe tracks and good jets in data (without electron-jet-overlap cut).

plicity samples. These corrections are applied as multiplicative factors to the measured jet multiplicities in data.




Figure 77. ΔR between probe tracks and good jets using PYTHIA MC (without electron-jet-overlap cut).

Jet Multiplicity	Electron-Jet-Overlap Coefficient
≥ 1	1.059 ± 0.028
≥ 2	1.075 ± 0.041
≥ 3	1.092 ± 0.054
≥ 4	1.109 ± 0.067
≥ 5	1.125 ± 0.077

TABLE XIII

ELECTRON-JET-OVERLAP COEFFICIENTS WITH SYSTEMATIC UNCERTAINTIES.




Figure 78. ΔR between probe tracks and good jets in data (after the electron-jet-overlap cut was applied).




Figure 79. ΔR between generated electrons ($p_T > 25$ GeV, $|\eta| < 1.1$) and partons ($p_T > 20$ GeV, $|\eta| < 2.5$).




Figure 80. ΔR between partons and matched calorimeter jets ($p_T > 20$ GeV, $|\eta| < 2.5$).

Jet Multiplicity	Total number of entries	Signal	Background
≥ 1	2,625.2	2,550.7 (52.5)	74.6
≥ 2	404.5	392.0 (8.1)	12.5
≥ 3	64.7	61.6 (1.3)	3.1
≥ 4	14.7	14.7 (-)	0
≥ 5	10.8	10.8 (-)	0

TABLE XIV

NUMBER OF SIGNAL EVENTS (DRELL-YAN IN PARENTHESIS) AND NUMBER OF BACKGROUND EVENTS (QCD) FOR DIFFERENT JET MULTIPLICITIES. NUMBERS ARE CORRECTED FOR TRIGGER, EM AND TRACKING INEFFICIENCIES (AFTER APPLYING THE UNSMEARING, JET RECO/ID AND ELECTRON-JET-OVERLAP CORRECTIONS).

9.2.3 Cross Sections

Figure 81 - Figure 85, shows the diem invariant mass distributions for jet multiplicities ≥ 1 to ≥ 5 which are used to extract the number of signal and background events for the cross section calculation (corrected for trigger, EM and tracking inefficiencies). For jet multiplicities of ≥ 1 and ≥ 2 , the same technique is followed as outlined in Chapter 8.2. For jet multiplicities of ≥ 3 , sidebands are used to estimate the background. For higher jet multiplicities there is not enough statistics to estimate any background.

Table XIV summarizes the number of signal and background events for each jet multiplicity. The fully corrected and unsmeared cross sections versus jet multiplicities (with jet $p_T > 20$ GeV, $|\eta| < 2.5$) are shown in Figure 86 with statistical errors.




Figure 81. Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 1$ jet sample. The solid line shows a Gaussian plus Breit-Wigner fit to the Z peak. The dashed line shows an exponential fit to the QCD and Drell-Yan contribution.




Figure 82. Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 2$ jet sample. The solid line shows a Gaussian plus Breit-Wigner fit to the Z peak. The dashed line shows an exponential fit to the QCD and Drell-Yan contribution.




Figure 83. Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 3$ jet sample.




Figure 84. Dielectron invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 4$ jet sample.




Figure 85. Diem invariant mass distribution for the $Z/\gamma^* \rightarrow e^+e^- + \geq 5$ jet sample.




Figure 86. $Z/\gamma^*(\rightarrow e^+e^-) + \geq n$ jet cross sections with statistical errors.

CHAPTER 10

SYSTEMATICS

This chapter evaluates various sources for systematic errors.

10.1 Jet Energy Scale Systematic Error

The error due to the jet energy scale (version 5.3) is estimated by varying the energy scale correction up and down by 1σ (combined systematic and statistical JES uncertainty) and subsequently recalculating the diem invariant mass histograms (corrected for trigger, EM reco/ID, and EM-Track matching inefficiencies) to get the number of corrected signal events for different jet multiplicities. After this step, the cross sections are recalculated to estimate the JES error. Figure 87 shows the effect of the JES error on the corrected jet multiplicity distribution.

The percentage errors for each jet multiplicity are fitted linearly to get a smooth curve and to be independent from statistical fluctuations. The final JES error is the maximum of the actual data error and the fit error. Table XV summarizes the JES errors. The JES errors are the dominant errors in this analysis.

10.2 Systematic Error of Cross Section Unfolding

A detailed description of the jet reco/ID efficiency errors can be found in Reference (70). Figure 88, Figure 89 and Figure 90 show the jet reco/ID efficiencies with error bands for central, ICR, and forward rapidities. The errors are due to the statistical uncertainty of the




Figure 87. $\pm 1\sigma$ fluctuation of the jet energy scale (JES 5.3)

Jet multiplicity	Cross Section	JES Errors
≥ 1	29.6 pb	± 2.9 pb
≥ 2	4.50 pb	± 0.83 pb
≥ 3	0.655 pb	± 0.185 pb
≥ 4	0.151 pb	± 0.057 pb
≥ 5	0.1035 pb	± 0.0520 pb

TABLE XV

FINAL CROSS SECTIONS WITH JET ENERGY SCALE ERRORS.

scale factor and straight efficiency, the missing E_T cut, and the differences between CKKW and PYTHIA MC generators. To estimate the jet reco/ID uncertainty, the unsmeared and jet reco/ID correction factors are rederived using the upper and lower error bands of the jet reco/ID efficiencies.

Figure 91 shows the scale factor used in this analysis overlaid with the average scale factor determined by the Top group. Both curves agree within the error bands.

As a cross check, a closure test is performed where the “efficiency” of finding a recoiling jet as a function of Z p_T is compared in data and particle level MC (Figure 92). Both the data and MC distributions have a jet p_T cut of 8 GeV applied. In addition, the MC particle jets are smeared with the data jet energy resolution. The data jet reco/ID efficiencies are also applied (by dropping jets) to the MC sample. The two distributions are in good agreement within errors. Table XVI summarizes the jet reco/ID errors.

The parameterization of the jet resolution is based on JES 5.0 with T42 applied (see Chapter

Jet multiplicity	Cross Section	Jet Reco/ID Errors
≥ 1	29.6 pb	+2.2 -1.6 pb
≥ 2	4.50 pb	+0.64 -0.57 pb
≥ 3	0.655 pb	+0.109 -0.105 pb
≥ 4	0.157 pb	+0.041 -0.028 pb
≥ 5	0.1035 pb	+0.0567 -0.0287 pb

TABLE XVI

FINAL CROSS SECTIONS WITH JET RECO/ID ERRORS.

7.2.1). The difference between the JES 5.0 and 5.3 parameterizations has to be taken into account. It has been shown (see Figure 93) that there is a change of approximately 5% between JES 5.1 and 5.3 (71). A conservative error of 10% is therefore assigned to account for the jet resolution uncertainty.

The error due to jet resolution smearing in the unsmearing procedure is estimated (Chapter 9.2.1) by varying the data jet resolution by $\pm 10\%$. Table XVII summarizes the jet resolution errors.

10.3 Electron-Jet-Overlap Systematic Error

For each jet multiplicity, electron-jet-overlap correction factors are derived using $\Delta R=0.4$ and $\Delta R=0.7$ rejection cones and then taking the middle value as the final correction. The systematic error is the difference between the middle values and the correction factors derived with $\Delta R=0.4$ and $\Delta R=0.7$. Additionally, a linear fit is applied to the final errors in order to




Figure 88. Jet reco/ID efficiencies with errors plotted versus particle jet p_T smeared with data energy resolution (central).




Figure 89. Jet reco/ID efficiencies with errors plotted versus particle jet p_T smeared with data energy resolution (ICR).




Figure 90. Jet reco/ID efficiencies with errors plotted versus particle jet p_T smeared with data energy resolution (forward).




Figure 91. Comparing scale factors between the Top group and this analysis. Equivalent techniques were used in both cases.




Figure 92. Comparing “efficiency” of finding a recoiling jet as a function of $Z p_T$ in data and MC.




Figure 93. Comparing jet resolution for JES 5.1 (upper curve) and JES 5.3 (lower curve) in the central region of the Calorimeter. The difference is approximately 5% over the whole range.

Jet Multiplicity	Cross Section	Jet Resolution Errors
≥ 1	29.6 pb	± 0.5 pb
≥ 2	4.50 pb	± 0.14 pb
≥ 3	0.655 pb	± 0.017 pb
≥ 4	0.151 pb	± 0.014 pb
≥ 5	0.1035 pb	± 0.0681 pb

TABLE XVII

FINAL CROSS SECTIONS WITH JET RESOLUTION ERRORS.

Jet Multiplicity	Cross Section	Electron-jet-overlap Errors
≥ 1	29.6 pb	± 0.8 pb
≥ 2	4.50 pb	± 0.17 pb
≥ 3	0.655 pb	± 0.032 pb
≥ 4	0.151 pb	± 0.009 pb
≥ 5	0.1035 pb	± 0.0071 pb

TABLE XVIII

FINAL CROSS SECTIONS WITH ELECTRON-JET-OVERLAP CUT UNCERTAINTIES.

compensate for statistical fluctuations.

Table XVIII summarizes the systematic errors for the electron-jet-overlap cut.

10.4 Luminosity Systematic Error

The error due to the uncertainty in the luminosity measurement is 6.5% (19). Table XIX summarizes the luminosity errors.

Jet multiplicity	Cross Section	Luminosity Errors
≥ 0	248.9 pb	± 16.2 pb
≥ 1	29.6 pb	± 1.9 pb
≥ 2	4.50 pb	± 0.29 pb
≥ 3	0.655 pb	± 0.043 pb
≥ 4	0.151 pb	± 0.010 pb
≥ 5	0.1035 pb	± 0.0067 pb

TABLE XIX

FINAL CROSS SECTIONS WITH LUMINOSITY ERRORS.

10.5 Systematic Errors Due to Efficiencies

In the following, the systematic uncertainties of the object based efficiencies are taken from Chapters 9.1.1 to 9.1.3, converted into event based systematic uncertainties, and then propagated to the cross sections.

10.5.1 Trigger Efficiency

A relative systematic error of $\pm 1\%$ is estimated due to the variations in the trigger efficiencies versus jet multiplicity (see Chapter 9.1.1):

$$\varepsilon_{Object}(\text{pre-v12, inclusive sample}) = 94.6\%, \quad \delta\varepsilon_{Object} = 5\% \quad (10.1)$$

$$\varepsilon_{Event} = 2 \cdot \varepsilon_{Object} - \varepsilon_{Object}^2 = 99.7\% \quad (10.2)$$

$$\varepsilon_{Event}(-1\sigma) = 2 \cdot (\varepsilon_{Object} - \delta\varepsilon_{Object}) - (\varepsilon_{Object} - \delta\varepsilon_{Object})^2 = 98.9\% \quad (10.3)$$

$$\text{Relative Error} = \frac{99.7\% - 98.9\%}{99.7\%} = 0.8\% \approx 1\% \quad (10.4)$$

10.5.2 EM Reconstruction and Identification Efficiency

A relative systematic error of $\pm 7\%$ is assumed due to the variations in the EM reco and ID efficiencies versus jet multiplicity (see Chapter 9.1.2):

$$\varepsilon_{Object}(\text{data, inclusive sample}) = 88.9\%, \quad \delta\varepsilon_{Object} = 3\% \quad (10.5)$$

$$\varepsilon_{Event} = \varepsilon_{Object}^2 = 79.0\% \quad (10.6)$$

$$\varepsilon_{Event}(-1\sigma) = (\varepsilon_{Object} - \delta\varepsilon_{Object})^2 = 73.8\% \quad (10.7)$$

$$\text{Relative Error} = \frac{79.0\% - 73.8\%}{79.0\%} = 6.6\% \approx 7\% \quad (10.8)$$

10.5.3 EM-Track Match Efficiency

Table XX summarizes the relative systematic uncertainties due to the variations in the EM-Track matching efficiencies versus jet multiplicity (see Chapter 9.1.3). The uncertainties are estimated based on the following equations:

$$\varepsilon_{Object}(\text{data, n-jet sample}) = \varepsilon_n, \quad \delta\varepsilon_{Object}(\text{data, n-jet sample}) = \delta\varepsilon_n \quad (10.9)$$

$$\varepsilon_{Event} = 2 \cdot \varepsilon_n - \varepsilon_n^2 \quad (10.10)$$

$$\varepsilon_{Event}(-1\sigma) = 2 \cdot (\varepsilon_n - \delta\varepsilon_n) - (\varepsilon_n - \delta\varepsilon_n)^2 \quad (10.11)$$

Jet Multiplicity	Relative Error
≥ 1	0.5%
≥ 2	1.5%
≥ 3	2.3%
≥ 4	3.7%
≥ 5	5.5%

TABLE XX

RELATIVE ERRORS DUE TO UNCERTAINTY IN EM-TRACK MATCHING EFFICIENCIES.

$$\text{Relative Error} = \frac{\varepsilon_{Event} - \varepsilon_{Event}(-1\sigma)}{\varepsilon_{Event}} \quad (10.12)$$

10.5.4 Overall Efficiency Systematic Error

Table XXI summarizes the overall systematic error due to the efficiencies after adding all contributions in quadrature.

10.6 Jet Promotion Systematic Error

The effect of additional jets from multiple interactions within the same beam crossing (*jet promotion*) is studied. Jet multiplicities of events that have exactly one reconstructed primary vertex are compared with events that have at least two reconstructed primary vertices (Table XXII).

The two samples are normalized with respect to the number of events in the inclusive jet multiplicity bin. Initially the single vertex sample contains 7,848 events and the 2 (or more)

Jet Multiplicity	Cross Section	Efficiency Errors
≥ 1	29.6 pb	± 2.1 pb
≥ 2	4.50 pb	± 0.32 pb
≥ 3	0.655 pb	± 0.047 pb
≥ 4	0.151 pb	± 0.011 pb
≥ 5	0.1035 pb	± 0.0093 pb

TABLE XXI

OVERALL SYSTEMATIC UNCERTAINTIES DUE TO EFFICIENCIES (TRIGGER, EM, TRACKING).

vertex sample contains 5,900 events.

The jet promotion effect is small since the discrepancy between the two samples is within the statistical error.

The average numbers of reconstructed primary vertices for different jet multiplicity samples are also compared (Table XXIII). No effect within statistical errors can be seen.

10.7 Statistical Uncertainty

The statistical uncertainty has the following contributions:

- The error due to the total number of corrected events δN_{corr} (corrected for Trigger, EM and Tracking inefficiencies) is estimated with the following equation:

$$\delta N_{corr} = \sqrt{(w_{average} \cdot \delta N_{uncorr})^2 + (N_{uncorr} \cdot \delta w_{average})^2} \quad (10.13)$$

Jet multiplicity	Exactly one primary vertex	At least two primary vertices
≥ 0	5,900	5,900
≥ 1	705	696
≥ 2	92	97
≥ 3	11	16
≥ 4	3	1
≥ 5	1	1

TABLE XXII

NUMBER OF EVENTS FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES WHEN REQUIRING EXACTLY ONE RECONSTRUCTED PRIMARY VERTEX AND AT LEAST TWO RECONSTRUCTED PRIMARY VERTICES. ENTRIES ARE NORMALIZED WITH RESPECT TO THE 2 VERTEX SAMPLE.

Jet multiplicity	Average number of primary vertices
≥ 1	1.583 ± 0.852
≥ 2	1.622 ± 0.911
≥ 3	1.733 ± 0.814
≥ 4	1.4 ± 0.8
≥ 5	2.0 ± 1.0

TABLE XXIII

AVERAGE NUMBER OF RECONSTRUCTED PRIMARY VERTICES FOR DIFFERENT JET MULTIPLICITIES.

where N_{corr} is the total number of corrected events, N_{uncorr} is the total number of uncorrected events, and $w_{average}$ is the average weight used to correct for EM, Trigger and Tracking inefficiencies (≈ 1.36).

- The error due to the number of background events δB is estimated by fitting an exponential function $a \cdot \exp(b \cdot x)$ to the measured number of QCD events (Figure 94) and then using the error due to the two fitting parameters a and b as the uncertainty. The error of the exponential fit takes into account that the fitting parameters a and b are correlated:

$$f(x) = a \cdot \exp(b \cdot x) \quad (10.14)$$

$$\delta f(x) = \sqrt{\left(\frac{\partial f}{\partial a} \cdot \delta a\right)^2 + \left(\frac{\partial f}{\partial b} \cdot \delta b\right)^2 + 2 \cdot \frac{\partial f}{\partial a} \cdot \frac{\partial f}{\partial b} \cdot \text{covariance}(a, b)}. \quad (10.15)$$

- An additional error due to the statistical uncertainty of the acceptances is added (see Table XI).
- The statistical uncertainty due to the unsmearing and jet reco/ID coefficients is added (see Chapter 9.2.1).

Table XXIV summarizes the statistical errors.




Figure 94. Exponential fit to the number of QCD events for different inclusive jet multiplicities.

Jet Multiplicity	Cross Section	Statistical Uncertainty
≥ 0	248.9 pb	± 2.5 pb
≥ 1	29.6 pb	± 0.8 pb
≥ 2	4.50 pb	± 0.32 pb
≥ 3	0.655 pb	± 0.129 pb
≥ 4	0.151 pb	± 0.070 pb
≥ 5	0.1035 pb	± 0.0738 pb

TABLE XXIV

CROSS SECTIONS WITH STATISTICAL ERRORS.

CHAPTER 11

CONCLUSIONS

11.1 Results

The $Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ jet cross sections for jet multiplicities of 0 to 5 have been measured. The results are presented in terms of absolute cross sections and cross section ratios normalized with respect to the inclusive cross section.

Table XXV summarizes all measured cross sections, together with uncertainties due to statistics, systematics, and luminosity. Also listed is the number of fully corrected signal events for each jet multiplicity sample. Table XXVI and Figure 95 compare the measured cross sections with results from MCFM and CKKW MC simulations (see Chapter 7.2). The CKKW cross sections are normalized with respect to the measured cross section for the 1-jet sample. The matrix element generation of the CKKW samples was done up to jet multiplicities of 3. Higher jet multiplicities are due to parton showering and hadronization simulated with PYTHIA.

Table XXVII and Figure 96 compare the cross section ratios with results from MCFM and CKKW MC simulations. For cross section ratios (R_n), the luminosity measurement uncertainties cancel. All other systematic uncertainties contribute as shown in the following equation:

$$\delta R_n = \frac{\delta \sigma_n}{\sigma_0}, \quad \text{with } \sigma_0 = 248.9 \text{ pb.} \quad (11.1)$$

Jet Multiplicity	# of Signal Events	$Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ Jet Cross Section
≥ 0	18,263.8	248.9 pb ± 2.5 (stat) ± 16.2 (lumi)
≥ 1	2,550.7	29.6 pb ± 0.81 (stat) $^{+4.3}_{-4.0}$ (sys) ± 1.9 (lumi)
≥ 2	391.9	4.50 pb ± 0.32 (stat) $^{+1.1}_{-1.1}$ (sys) ± 0.29 (lumi)
≥ 3	63.6	0.655 pb ± 0.13 (stat) $^{+0.22}_{-0.22}$ (sys) ± 0.043 (lumi)
≥ 4	15.3	0.151 pb ± 0.070 (stat) $^{+0.072}_{-0.066}$ (sys) ± 0.010 (lumi)
≥ 5	6.8	0.104 pb ± 0.074 (stat) $^{+0.10}_{-0.06}$ (sys) ± 0.0067 (lumi)

TABLE XXV

CROSS SECTIONS FOR DIFFERENT INCLUSIVE JET MULTIPLICITIES. NUMBER OF SIGNAL EVENT ENTRIES HAVE UNSMEARING, JET RECO/ID AND ELECTRON-JET-OVERLAP CORRECTIONS APPLIED.

Jet p_T distributions for different jet multiplicities are compared between data and ALPGEN+PYTHIA MC simulations (Figure 97).

The results are in good agreement with QCD predictions.

11.2 Future Plans

Future work on this analysis at DØ will include the use of CKKW samples with matrix element generation of up to 5 jets. This will further improve cross section comparisons at higher jet multiplicities, as well as unsmeared results.

Additional background processes also need to be evaluated. Preliminary studies indicate that contributions from backgrounds such as $Z/\gamma^* \rightarrow \tau^+\tau^-$ are small.

Jet Multiplicity	Measured Cross Section	MCFM	CKKW
≥ 0	248.9 pb ± 16.4 (tot)	241.5 pb ± 0.1 (stat)	-
≥ 1	29.6 pb $^{+4.8}_{-4.6}$ (tot)	26.2 pb ± 0.044 (stat)	29.6 pb
≥ 2	4.50 pb $^{+1.2}_{-1.2}$ (tot)	5.21 pb ± 0.069 (stat)	5.22 pb
≥ 3	0.655 pb $^{+0.26}_{-0.26}$ (tot)	-	0.798 pb
≥ 4	0.151 pb $^{+0.10}_{-0.10}$ (tot)	-	0.096 pb
≥ 5	0.104 pb $^{+0.13}_{-0.10}$ (tot)	-	0.008 pb

TABLE XXVI

COMPARISON OF MEASURED CROSS SECTIONS WITH RESULTS FROM MCFM AND CKKW.

Jet multiplicity	$\frac{\sigma(Z/\gamma^*(\rightarrow e^+e^-)+\geq n Jets)}{\sigma_{Z/\gamma^*}} [\cdot 10^{-3}]$	MCFM	CKKW
≥ 1	119.1 ± 3.3 (stat) $^{+17.2}_{-16.2}$ (sys)	108.4	119.1
≥ 2	18.1 ± 1.3 (stat) $^{+4.5}_{-4.3}$ (sys)	21.6	21.0
≥ 3	2.6 ± 0.52 (stat) $^{+0.90}_{-0.89}$ (sys)	-	3.2
≥ 4	0.61 ± 0.28 (stat) $^{+0.29}_{-0.27}$ (sys)	-	0.39
≥ 5	0.42 ± 0.30 (stat) $^{+0.42}_{-0.24}$ (sys)	-	0.03

TABLE XXVII

COMPARISON OF MEASURED CROSS SECTION RATIOS WITH RESULTS FROM MCFM AND CKKW.




Figure 95. $Z/\gamma^*(\rightarrow e^+e^-)+\geq n$ jet cross sections in data (with total errors) compared with MCFM and CKKW.




Figure 96. Ratios of the $Z/\gamma^*(\rightarrow e^+e^-) + \geq n$ jet cross sections to the total inclusive $Z/\gamma^* \rightarrow e^+e^-$ cross section versus n . The errors on the data include the combined statistical and systematic uncertainties. The dashed line (CKKW) represents the predictions of LO matrix element calculations using PYTHIA for parton showering and hadronization, normalized to the measured $Z/\gamma^* + \geq 1$ jet cross section ratio. The diamonds represent the MCFM predictions.




Figure 97. Data to theory (ALPGEN+PYTHIA) comparison for the highest p_T jet distribution in the $Z + \geq 1$ jet sample, for the second highest p_T jet distribution in the $Z + \geq 2$ jet sample and for the third highest p_T jet distribution in the $Z + \geq 3$ jet sample. The errors on the data are only statistical.

APPENDICES

Appendix A

THE DØ COLLABORATION

R. Piegaia, A. Salles

Universidad de Buenos Aires, Buenos Aires, Argentina

G.A. Alves, J. Barreto, H. da Motta, M.-E. Pol, M.S. Rangel, F.A.S. Rezende, M. Souza,

M. Vaz

LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

A.C.S. Assis Jesus, M. Begalli, W. Carvalho, C. De Oliveira Martins, S. Greder,

J.R.P. Mahon, H.B. Malbouisson, J. Molina, L. Mundim, V. Oguri, N. Oliveira,

W.L. Prado da Silva, R.F. Rodrigues, A. Santoro, A. Sznajder

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

E.M. Gregores, S.M. Lietti, P.G. Mercadante, S.F. Novaes

Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

S. Beale, C. Belanger-Champagne, K. Chan, Y. Coadou, D. Gillberg, Z. Liu, R.W. Moore,

D.C. O'Neil, W. Taylor, B. Vachon, M.C. Vetterli

University of Alberta, McGill University, Simon Fraser University and York

University, Canada

H.S. Mao

Institute of High Energy Physics, Beijing, People's Republic of China

L. Han, J. Yi

University of Science and Technology of China, Hefei, People's Republic of China

C. Avila, B. Gómez, D. Mendoza, J.P. Negret, R. Ramirez, J.M.R. Roldan

Universidad de los Andes, Bogotá, Colombia

V. Hynek, J. Kvita, R. Leitner, K. Soustruznik

Charles University, Center for Particle Physics, Prague, Czech Republic

P. Homola, Z. Hubacek, R. Otec, V. Simak, P. Vokac

Czech Technical University, Prague, Czech Republic

A. Kupco, M. Lokajicek, V. Vrba

**Institute of Physics, Academy of Sciences, Center for Particle Physics, Prague,
Czech Republic**

B. Hoeneisen

Universidad San Francisco de Quito, Quito, Ecuador

F. Badaud, P. Gay, Ph. Gris, V. Lesne, F. Tissandier

**Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal,
Clermont-Ferrand, France**

Y. Arnoud, F. Chevallier, S. Crépe-Renaudin, A.-M. Magnan, G. Sajot, J. Stark, C. Yu

**Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS,
Université de Grenoble, Grenoble, France**

L. Berntzon, S. Calvet, M.-C. Cousinou, A. Duperrin, E. Kajfasz, S. Kermiche, A. Mendes,
E. Nagy, M. Talby

CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France

L. Duflot, J.-F. Grivaz, M. Jaffré, N. Makovec, P. Pétroff

Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS, Orsay, France

B. Andrieu, U. Bassler, G. Bernardi, E. Busato, L. Sonnenschein, S. Trincas-Duvoid,
J.-R. Vlimant

LPNHE, Universités Paris VI and VII, IN2P3-CNRS, Paris, France

M. Agelou, M. Besançon, F. Déliot, P. Demine, P. Lutz, M. Michaut, E. Perez, C. Royon,
V. Shary, B. Tuchming, D. Vilanova

DAPNIA/Service de Physique des Particules, CEA, Saclay, France

J.-L. Agram, D. Bloch, F. Charles, B. Clément, A. Gay, W. Geist, D. Gelé, A.-C. Le Bihan,
A. Lounis, I. Ripp-Baudot, V. Siccardi

**IReS, IN2P3-CNRS, Université Louis Pasteur Strasbourg, and Université de
Haute Alsace, France**

N. Estre, G. Grenier, T. Kurča, P. Lebrun, J.P. Martin, T. Millet, G.S. Muanza, P. Verdier
**Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude
Bernard, Villeurbanne, France**

C. Autermann, M. Erdmann, T. Hebbeker, D. Käfer, C. Magass, A. Meyer
RWTH Aachen, III. Physikalisches Institut A, Aachen, Germany

J. Meyer, A. Quadt, C. Schwanenberger, N. Wermes
Universität Bonn, Physikalisches Institut, Bonn, Germany

U. Blumenschein, V. Buescher, I. Fleck, H. Fox, G. Herten, K. Jakobs, J.-P. Konrath,
O. Mundal, C. Noeding, M. Titov, I. Torchiani

Universität Freiburg, Physikalisches Institut, Freiburg, Germany

C. Ay, M. Hohlfeld, T. Kuhl, D. Meder, S. Tapprogge, T. Trefzger, C. Zeitnitz
Universität Mainz, Institut für Physik, Mainz, Germany

O. Biebel, M. Binder, J. Elmsheuser, F. Fiedler, P. Haefner, B. Leonhardt, T. Nunnemann,
D. Schaile, P. Schieferdecker, R. Ströhmer

Ludwig-Maximilians-Universität München, München, Germany

K. Hamacher, A. Harel, H. Hoeth, S. Kersten, P. Mättig, C. Schmitt, M. Vaupel

Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

S.B. Beri, V. Bhatnagar, R. Kaur, J.M. Kohli

Panjab University, Chandigarh, India

B. Choudhary, A. Kumar, M. Naimuddin, K. Ranjan, R.K. Shivpuri

Delhi University, Delhi, India

B.S. Acharya, P. Banerjee, S. Banerjee, S. Chakrabarti, A. Chandra, S.R. Dugad, P.K. Mal,

N.K. Mondal, K.J. Rani

Tata Institute of Fundamental Research, Mumbai, India

M. Cwiok, M.W. Grünewald

University College Dublin, Dublin, Ireland

S.H. Ahn, S.J. Hong, T.J. Kim, K.S. Lee, S.K. Park

Korea Detector Laboratory, Korea University, Seoul, Korea

H. Castilla-Valdez, E. De La Cruz-Burelo, P.L.M. Podesta-Lerma, A. Sánchez-Hernández

CINVESTAV, Mexico City, Mexico

K. Bos, S. Caron, P. de Jong, J.G. Hegeman, P. Houben, G.G.G. Massaro, P.J. van den Berg,

W.M. van Leeuwen, M. Vreeswijk

**FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam,
The Netherlands**

M. Anastasoae, L.S. Ancu, S.J. de Jong, F. Filthaut, C.F. Galea, M.H. Kirby, N.A. Naumann

Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands

V.M. Abazov, G.D. Alexeev, D.V. Bandurin, G. Erusalimtshev, G. Golovanov, A.M. Kalinin,

Y.M. Kharzheev, E.V. Komissarov, V.L. Malyshev, Y.P. Merekov, S.Y. Porokhovoi,

N.A. Russakovich, A.A. Shishkin, N.B. Skachkov, V.V. Tokmenin, L.S. Vertogradov,

Y.A. Yatsunenko

Joint Institute for Nuclear Research, Dubna, Russia

A. Evdokimov, V. Gavrilov, P. Polozov, G. Safronov, V. Stolin, V.I. Turtikov

Institute for Theoretical and Experimental Physics, Moscow, Russia

E.E. Boos, S. Bunichev, L.V. Dudko, P. Ermolov, D. Karmanov, A. Koubarovsky,

V.A. Kuzmin, A. Leflat, M. Merkin, V.I. Rud, S. Sherstnev, E.G. Zverev

Moscow State University, Moscow, Russia

V.A. Bezzubov, S.P. Denisov, S.A. Efremov, V.N. Evdokimov, A.V. Ferapontov,
A.K. Klimenko, V.M. Korablev, V.I. Koreshev, A.V. Kozelov, E.A. Kozlovsky, V.V. Lipaev,
A.V. Popov, A.A. Shchukin, Y.V. Sinkin, D.A. Stoyanova, I.A. Vasilyev, S.A. Zvyagintsev

Institute for High Energy Physics, Protvino, Russia

G. Alkhazov, S. Anufriev, V. Kim, A. Lobodenko, P. Neustroev, G. Obrant, Y. Scheglov,

L. Uvarov, S. Uvarov

Petersburg Nuclear Physics Institute, St. Petersburg, Russia

B. Åsman, C. Clément, P. Eerola, T. Ekelöf, N. Gollub, P. Hansson, S. Lager,

B. Lund-Jensen, T. Moa, J. Strandberg

Lund University, Royal Institute of Technology, Stockholm University, and

Uppsala University, Sweden

M. Audsley Hammond, I. Bertram, C. Biscarat, G. Borissov, B. Davies, M. Doidge,

A.J. Finch, M. Lewin, P. Love, A. Rakitine, P.N. Ratoff, A. Sopczak

Lancaster University, Lancaster, United Kingdom

C. Barnes, R. Beuselinck, I. Blackler, F. Blekman, C.P. Buszello, G. Davies, J.F. Hassard,

A. Jenkins, R. Jesik, P. Jonsson, P. Lewis, L. Lobo, M. Petteni, S. Robinson, T. Scanlon,

F. Villeneuve-Seguiér, P. Vint

Imperial College, London, United Kingdom

B. Cox, T. Edwards, M. Ford, J. Monk, M. Owen, K. Peters, S. Söldner-Rembold, P. Telford,

A.S. Turcot, T.R. Wyatt

University of Manchester, Manchester, United Kingdom

N.M. Giao, D. Han, T.M. Ngoc, H.T.K. Trang, T.D. Tu

Hochiminh City Institute of Physics, Hochiminh City, Vietnam

S. Anderson, S. Burke, E. Cheu, B. Gmyrek, K. Johns, J. Leveque, R. McCroskey, M. Shupe,

P. Tamburello, J. Temple, E.W. Varnes

University of Arizona, Tucson, Arizona 85721, USA

R.J. Madaras, M. Strovink

Lawrence Berkeley National Laboratory and University of California, Berkeley,

California 94720, USA

R.E. Hall

California State University, Fresno, California 93740, USA

S. Choi, R. Clare, J. Ellison, R. Gelhaus, A.P. Heinson, I. Iashvili, P.M. Perea, S.J. Wimpenny

University of California, Riverside, California 92521, USA

T. Adams, A. Askew, S. Blessing, N.J. Buchanan, D. Duggan, Y. Gershtein, S. Hagopian,
V. Hagopian, D. Kau, J. Lazoflores, W.M. Lee, J. Miao, S. Nelson, H.B. Prosper, J. Sekaric,
S. Sengupta, S. Sumowidagdo, S. Tentindo-Repond, H.D. Wahl

Florida State University, Tallahassee, Florida 32306, USA

B. Baldin, J.F. Bartlett, L. Bellantoni, R. Bernhard, P.C. Bhat, A. Boehnlein, A. Bross,
S. Burdin, W.E. Cooper, M. Demarteau, D. Denisov, H.T. Diehl, M. Diesburg, V.D. Elvira,
J. Estrada, H.E. Fisk, S. Fu, S. Fuess, E. Gallas, K. Gounder, H. Greenlee, S. Grünendahl,
G. Gutierrez, K. Hanagaki, R. Illingworth, A.S. Ito, M. Johnson, A. Jonckheere, A. Juste,
P. Kasper, B. Klima, S. Krzywdzinski, F. Lehner, Q.Z. Li, D. Lincoln, R. Lipton, A.L. Lyon,
Y. Maravin, M. Martens, K.W. Merritt, M. Mulders, A. Nomerotski, V. O'Dell, N. Oshima,
V.M. Podstavkov, P.A. Rapidis, P. Rubinov, G. Savage, V. Sirotenko, R.P. Smith, L. Stutte,
M. Tomoto, M. Weber, D. Wicke, M. Wobisch, J. Womersley, R. Yamada, T. Yasuda,
M. Zanabria, D. Zhang

Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

M. Adams, M. Buehler, M. Camuyrano, C.E. Gerber, J.M. Heinmiller, G.J. Otero y Garzón,
E. Shabalina, A. Stone, T. Ten, N. Varelas

University of Illinois at Chicago, Chicago, Illinois 60607, USA

L. Bagby, G. Blazey, D. Chakraborty, A. Dyshkant, M. Eads, M. Fortner, D. Hedin,
J.G.R. Lima, A.K.A. Maciel, X. Song, N.M. Tran, S. Uzunyan, A. Zatserklyaniy, V. Zutshi

Northern Illinois University, DeKalb, Illinois 60115, USA

T. Andeen, D. Buchholz, G.A. Davis, M. Gagliardi, J. Hays, H. Schellman, D. Strom,
S. Yacoob, S.W. Youn

Northwestern University, Evanston, Illinois 60208, USA

R. Abrams, D. Bauer, J. Rieger, K. Stevenson, R. Van Kooten, L. Welty, D. Zieminska,
A. Zieminski

Indiana University, Bloomington, Indiana 47405, USA

O. Boeriu, N.M. Cason, E. Galyaev, A. Goussiou, M.D. Hildreth, D. Karmgard,
A. Kharchilava, D. Lam, H. Luo, M. Lynker, Y. Pogorelov, R. Ruchti, W.D. Shephard,
P. Svoisky, J. Torborg, J. Warchol, M. Wayne, N. Xuan

University of Notre Dame, Notre Dame, Indiana 46556, USA

O. Atramentov, J.M. Hauptman

Iowa State University, Ames, Iowa 50011, USA

P. Baringer, A. Bean, L. Christofek, D. Coppage, J. Gardner, C. Hensel, S. Jabeen,
T. Moulik, G.W. Wilson

University of Kansas, Lawrence, Kansas 66045, USA

M. Ahsan, T.A. Bolton, K. Harder, D. Onoprienko, F. Rizatdinova, M. Shamim, R.A. Sidwell,

M. Smith, E. Von Toerne

Kansas State University, Manhattan, Kansas 66506, USA

M. Das, Z.D. Greenwood, N. Parashar, L. Sawyer, J. Steele

Louisiana Tech University, Ruston, Louisiana 71272, USA

A. Baden, S. Eno, N.J. Hadley, C. Jarvis, S. Kunori, J. Mans, M.P. Sanders, T. Toole,

M. Verzocchi, L. Wang, M. Wetstein, M. Yan

University of Maryland, College Park, Maryland 20742, USA

K.M. Black, J.M. Butler, D.K. Cho, A. Das, S.N. Fatakia, L. Feligioni, U. Heintz, J. Kasper,

M. Narain

Boston University, Boston, Massachusetts 02215, USA

G. Alverson, E. Barberis, S. Doulas, R. Harrington, G. Hesketh, S. Reucroft, D. Shpakov,

D.R. Wood

Northeastern University, Boston, Massachusetts 02115, USA

A. Alton, J.D. Degenhardt, A. Magerkurth, H.A. Neal, J. Qian, Z. Zhao, B. Zhou

University of Michigan, Ann Arbor, Michigan 48109, USA

M. Abolins, J. Benitez, R. Brock, J. Dyer, D. Edmunds, R. Hauser, J. Kalk, J. Kozminski,
J. Linnemann, B.G. Pope, R. Schwienhorst, R. Unalan, H. Weerts

Michigan State University, East Lansing, Michigan 48824, USA

A. Melnitchouk, B. Quinn

University of Mississippi, University, Mississippi 38677, USA

A. Bellavance, K. Bloom, D. Claes, A. Dominguez, C. Lundstedt, S. Malik, G.R. Snow,
M. Voutilainen

University of Nebraska, Lincoln, Nebraska 68588, USA

W. Fisher, J. Haley, A. Schwartzman, C. Tully

Princeton University, Princeton, New Jersey 08544, USA

T. Bose, G. Brooijmans, H. Evans, A. Haas, C. Johnson, I. Katsanos, D. Khatidze,
B. Kothari, S. Lammers, J. Mitrevski, M. Mulhearn, J. Parsons, P.M. Tuts

Columbia University, New York, New York 10027, USA

M. Begel, J. Cammin, K.M. Chan, R. Demina, T. Ferbel, C. Garcia, G. Ginther, A. Khanov,
T. Negrini, S.-J. Park, M.-A. Pleier, P. Slattery, M. Zielinski

University of Rochester, Rochester, New York 14627, USA

S. Desai, H. Dong, P.D. Grannis, J. Guo, J.D. Hobbs, Y. Hu, R. McCarthy, Y.D. Mutaf,
N. Parua, M. Rijssenbeek, R.D. Schamberger, S. Towers, D. Tsybychev, A. Yurkewicz, J. Zhu

State University of New York, Stony Brook, New York 11794, USA

S. Kahn, J. Kotcher, A. Patwa, S. Protopopescu, S. Snyder, K. Yip

Brookhaven National Laboratory, Upton, New York 11973, USA

J. Snow

Langston University, Langston, Oklahoma 73050, USA

B. Abbott, P. Gutierrez, I. Hall, S. Jain, M. Kopal, A. Pompoš, H. Severini, P. Skubic,

M. Strauss

University of Oklahoma, Norman, Oklahoma 73019, USA

B.C.K. Casey, D. Chapin, D. Cutts, R. Hooper, S. Kesisoglou, G. Landsberg,

S.E.K. Mattingly, R. Partridge, Y. Xie, H.D. Yoo

Brown University, Providence, Rhode Island 02912, USA

A. Brandt, D. Brown, K. De, F. Jaafari, V. Kaushik, H. Kim, J. Li, M. Sosebee, B. Spurlock,

M.A. Strang, A. Vartapetian, A. White, J. Yu

University of Texas, Arlington, Texas 76019, USA

R. Kehoe, P. Renkel

Southern Methodist University, Dallas, Texas 75275, USA

P. Bargassa, M. Cooke, M. Corcoran, S.J. Lee, D. Mackin, H. Miettinen, P. Padley,

G. Pawloski

Rice University, Houston, Texas 77005, USA

R. Hirosky, A. Kryemadhi, K. Nelson

University of Virginia, Charlottesville, Virginia 22901, USA

T.H. Burnett, T. Gadfort, A. Garcia-Bellido, G. Gaudio, H.J. Lubatti, G. Watts, T. Zhao

University of Washington, Seattle, Washington 98195, USA

Appendix B

THE LEVEL 2 TRIGGER SYSTEM AND ALPHA BOARDS

Level 2 is an essential part of the DØ trigger and data acquisition system (72; 73; 74; 75). It was designed to provide a reduction in the event rate by a factor of 10 within a 100 μsec time window while inducing less than 5% deadtime. As outlined in Chapter 4.2 the L2 trigger system is organized as a two-stage *stochastic pipeline*. In the first step (*preprocessing stage*) sub-detector based proto-objects are formed by preprocessors. The second step (*global processing*) combines the information provided by the preprocessors to make the event-wide L2 trigger decision. One advantage of this design is that each preprocessor is able to operate in parallel, independently from all other preprocessors. This avoids deadtime since decision times can vary significantly from event to event. Additionally up to 16 events can be queued in buffers (*FIFOs*) between the stages. Figure 98 illustrates the relation between sub-detectors, and L1/L2 trigger elements.

B.1 The Standard Level 2 Crate

All L2 systems occupy 9U¹ VME crates (76):

- Forward Muon Preprocessor (L2MUF)
- Central Muon Preprocessor (L2MUC)

¹1U = 1.75 in




Figure 98. Schematic view of subdetectors with L1 and L2 trigger elements. Horizontal arrows indicate the direction of dataflow.

- Calorimeter Preprocessor (L2CAL)
- Preshower Preprocessor (L2PS)
- Tracking Preprocessor (L2CTT/L2STT)
- Global Processor (L2GBL)

Each crate contains a 64-bit VME bus (77) and a custom-built high-speed bus (*Magic Bus*, or MBus) for data-handling on an auxiliary backplane (78). The 128 bit MBus supports data rates of up to 320 Mbit/s. Figure 99 shows a schematic view of the L2CAL crate with its main components:

MPM (*Multi Port Memory*): A Bit3 MPM card is used as the VME crate controller. It downloads run-time parameters and reports monitoring data to the Trigger Control Computer (TCC¹).

VBD (*VME Buffer Driver*): A VBD card is used to write data to the L3 system. The VBD cards were later replaced by VME single board computers (SBC).

Workers/Administrator (*Alpha Boards*): The Alpha Boards perform the main processing step within the L2 trigger system. Their main purpose is the application of the L2 trigger algorithms (*Worker*). They also handle additional event processing and local trigger control tasks (*Administrator*). A detailed overview of the Alpha Board system architecture is given in Chapter B.2. The Alpha Boards were later substituted by *Beta Boards* (79).

¹TCC handles run control, downloads run-specific information, and collects monitoring data.

MBT (*Magic Bus Transceiver*): The MBT cards (80) receive L1 information and broadcast it to the Worker/Administrator via the MBus. After preprocessing is finished, the outputs are sent to the L2 Global Processor (L2GBL). The MBT cards also receive information regarding L1 trigger accepts, L2 trigger decisions, and system-wide initializations (*Serial Command Link Initialize*, or SCLinit). Depending on the processing load, additional *Assistant MBTs* can be added to the *Pilot MBT*.

Specialized hardware: In some cases additional specialized hardware for data conversion and processing is added. In the case of the L2CAL crate, the L1 input signals arrive in the form of fiber-optic cables. A special converter card (*Fiber Input Converter*, or FIC) translates optical signals into electrical signals (81).

B.2 Level 2 Alpha Processors

The overall design of the Alpha Board (Figure 100, Reference (82)) is based on the layout of the DEC ¹ PC164 motherboard (83). It features a 500 MHz Alpha CPU running under real-time Linux. In addition to the internal PCI bus several elements supporting VME and MBus interfaces have been added. Figure 101 shows a schematic view of the Alpha Board. The main components are described in the following.

B.2.1 PC164 Based Design

The Alpha Board uses the first commercially available 64-bit RISC processor: a 500 MHz Alpha 21164 CPU (84; 85). It can execute 2-4 instructions per cycle. The 500 MHz CPU

¹Digital Equipment Corp. merged with Compaq Computer Corp. in 1998.




Figure 99. Layout of Level 2 calorimeter (L2CAL) crate with main components.

frequency is generated by a 50 MHz oscillator by means of a divide-by-10 phase-locked loop circuit.

The Alpha Board has several caches implemented, both on-chip (integrated into the CPU) and external. 8 kB data and instruction caches (Dcache and Icache, respectively) buffer the most frequently used data and instructions to speed up processing. A secondary on-chip 96 kB mixed data and instruction cache (Scache) is also used. A third level of external cache (4 MB) is not being utilized.

An interface between the CPU, main memory, and the PCI bus is provided by the 21172 Core Logic Chipset (Figure 102, Reference (86)). The 21172 Chipset consists of the 21172-CA chip (CIA) which provides the interface between the CPU and the PCI bus. It also provides




Figure 100. Level 2 Alpha Board with Ethernet card and hard disk drive.




Figure 101. Block diagram of Level 2 Alpha Board.

main memory controlling and addressing capabilities. A set of four data switching chips (DSW) functions as a multiplexing/demultiplexing device for main memory access. The 21172 Chipset provides data transfer rate of up to 267 MB/s (64-bit at 30 ns/cycle).

The Alpha Board carries two kinds of firmware. A 128 kB SROM chip is directly connected to the Icache. It contains the boot code that is loaded into the CPU at power-up, including the initialization of the PCI bus, the setting of registers in the CIA, the setting up of the various caches and memory, and finally the copying of the contents of the Flash ROM to memory. The 1 MB Flash ROM chip contain OS and task specific instructions. This includes instructions to download user code via a PCI Ethernet card, a rudimentary local debugger ("Debug Monitor"), and the server for a remote debugger. The SROM chip is installed in a socket, and can therefore

be programmed by using an external device programmer. The Flash ROM chip is soldered to the board, and must be programmed in situ.

A bank of eight SIMM slots provides 128 MB of DRAM memory.

The 64-bit 33 MHz PCI bus has two expansion slots. A standard PCI Ethernet card connected to one of the PCI slots is used for remote access. Two programmable chips (PLDs) are used for PCI interrupt handling and bus arbitration.

A PCI-IDE interface chip connects a 6 GB IDE hard disk drive, which has real-time Linux installed.

A PCI-ISA bridge provides support for external devices such as a mouse, keyboard, and floppy drive. It also supports a parallel port, two serial ports, and a real time clock. The ISA bus is used to access the Flash ROM.

B.2.2 Magic Bus Programmed Input/Output

An interface between MBus and PCI bus is provided by the Magic Bus Programmed I/O chip (MBusPIO). It reads and writes data between the two address spaces and thereby provides a way of communication between different cards that are connected to the MBus. For example, a board connected to the MBus can write or read into Alpha Board memory. A programmable Xilinx XC4036EX chip (MAGICFPGA) is used for this task.

B.2.3 Magic Bus Direct Memory Access Interface

In order to boost processing time, data is transferred to and from Alpha Board memory by direct memory access (DMA), i.e. data on the MBus goes directly to Alpha Board memory




Figure 102. Block diagram of the 21172 Core Logic Chipset.

without involving the CPU. This *DMA engine* is implemented in two programmable chips: a Xilinx FPGA, and a Cypress CPLD.

B.2.4 VME Interface

A 64-bit PCI-VME interface is provided by the Tundra Universe II chip by mapping “windows” of VME address space to Alpha Board memory space (87).

B.2.5 TSI Interface

The TSI interface is used to receive and send additional information to the trigger system. Is is also used to monitor the state of the Alpha Board processing elements.

CITED LITERATURE

1. Arnison, G. et al.: Experimental observation of lepton pairs of invariant mass around 95 GeV/c² at the CERN SPS collider. Phys. Lett., 126B:398, 1983.
2. Bagnaia, P.: Evidence for Z(e+e-) at the CERN ppbar collider. Phys. Lett., 129B:130, 1983.
3. Veltman, M.: Facts and Mysteries in Elementary Particle Physics. World Scientific, 2003.
4. Griffiths, D.: Introduction to Elementary Particles. John Wiley & Sons, 1987.
5. Halzen, F. and Martin, A.: Quarks and Leptons. John Wiley & Sons, 1984.
6. Quigg, C.: Gauge Theories of the Strong, Weak, and Electromagnetic Interactions. Westview Press, 1983.
7. Ellis, R. et al.: QCD and Collider Physics. Cambridge University Press, 1996.
8. Prosper, H. and Danilov, M.: Techniques and Concepts of High-Energy Physics XII. NATO Science Series, 2003.
9. Nurse, E.: A Measurement of the Inclusive $Z/\gamma^* \rightarrow \mu^+\mu^-$ Cross Section and Study of W and Z Events in $p\bar{p}$ Collisions at D0. University of Manchester, 2005.
10. The Particle Adventure. 2002.
<http://particleadventure.org/particleadventure/>.
11. CDF: Properties of Jets in Z Boson Events from 1.8 TeV p-barp Collisions. Physical Review Letters, 77:448, 1996.
12. Berends et al.: Exact expressions for processes involving a vector boson and up to five partons. Nuclear Physics B, 321:39–82, 1989.
13. Thompson, J.: Introduction to Colliding Beams at Fermilab. FERMILAB-TM-1909, 1994.

14. General Reference Documents for Accelerators. 2002.
http://www-numi.fnal.gov/workgrps/protonwg/accel_reference.html.
15. Accelerator Concepts Rookie Books. 2003.
http://www-bdnew.fnal.gov/operations/rookie_books/rbooks.html.
16. Mutař, Y.: Measurement of the Ratio of Inclusive Cross Sections $\sigma(p\bar{p} \rightarrow Z + b\text{-jet})/\sigma(p\bar{p} \rightarrow Z + \text{jet})$ at $\sqrt{s} = 1.96$ TeV. Stony Brook University, 2005.
17. Fermilab Anti-Proton Source Rookie Book. 1999.
http://www-bdnew.fnal.gov/pbar/documents/PBAR_Rookie_Book.PDF.
18. Edmunds, D.: Run IIa Tevatron Beam Structure.
http://www.pa.msu.edu/hep/d0/ftp/l1/framework/drawings/run_ii_a_beam_structure.ps.
19. Edwards, T. et al.: Determination of the Effective Inelastic p anti-p Cross Section for the D0 Run II Luminosity Measurement. FERMILAB-TM-2278-E, 2004.
20. Begel, M. et al.: D0 Luminosity in Run 2: Delivered. D0 Note 3970, 2002.
21. Ferbel, T.: A Brief Description of the D0 Detector in Run II. 2004.
<http://www-d0.fnal.gov/Run2Physics/WWW/templates/detector.tex>.
22. D0 Collaboration: The Upgraded D0 Detector. Preprint submitted to Elsevier Science, 2005.
23. D0 Collaboration: D0 Silicon Tracker Technical Design Report. 1994.
http://d0server1.fnal.gov/projects/silicon/www/tdr_final.ps.
24. D0 Collaboration: The D0 Upgrade: Central Tracker Technical Design Report. 1999.
http://d0server1.fnal.gov/users/stefan/www/CFT_TDR/CFT_TDR.ps.
25. Yurkewicz, A.: Search for Evidence of Supersymmetry in the Like-sign Dimuon Channel at the D0 Experiment. Michigan State University, 2004.
26. Adams, M. et al.: Design Report of the Central Preshower Detector for the D0 Upgrade. 1996.
<http://d0server1.fnal.gov/users/qianj/CPS/doc/dn3104.pdf>.

27. Gordeev, A. et al.: Technical Design Report of the Forward Preshower Detector for the D0 Upgrade. D0 Note 3445, 1998.
28. Abachi, S. et al.: The D0 Detector. Nucl. Instrum. Methods Phys. Res. A, 338:185, 1994.
29. Snyder, S.: Measurement of the Top Quark Mass at D0. Stony Brook University, 1995.
30. Baldin, B. et al.: Technical Design Report of the Central Muon System. D0 Note 3365, 1997.
31. Alexeev, G. et al.: Technical Design Report of the D0 Forward Muon Tracking Detector Based on Mini-Drift Tubes. D0 Note 3366, 1997.
32. Blazey, J.: The D0 Run II Trigger.
<http://niuhep.physics.niu.edu/~blazey/rt.ps>.
33. D0 Collaboration: d0reco. 2004.
<http://www.d0.fnal.gov/Run2Physics/WWW/algorithm.htm>.
34. Borisov, G.: Ordering a Chaos... or Technical Details of AA Tracking. 2004.
http://www-d0.fnal.gov/global_tracking/talks/20030228/talk-adm-030228.ps.
35. Khanov, A.: HTF: Histogramming Method for Finding Tracks. D0 Note 3778, 2000.
36. Haas, A.: A Search for Neutral Higgs Bosons at High $\tan\beta$ in Multi-jet Events from $p\bar{p}$ Collisions at $\sqrt{s} = 1960$ GeV. University of Washington, 2004.
37. Schwartzman, A. and Narain, M.: Probabilistic Primary Vertex Selection. D0 Note 4042, 2002.
38. Melnitchouk, A.: Search for non-SM Light Higgs Boson in the $h \rightarrow \gamma\gamma$ Channel. Brown University, 2004.
39. Blazey et al.: Run II Jet Physics. hep-ex/0005012, 2000.
40. Busato, E. and Andrieu, B.: Jet Algorithms in the D0 Run II Software: Description and User's Guide. D0 Note 4457, 2004.
41. Agram, J.-L. et al.: Jet Energy Scale at D0 Run II. D0 Note 4720, 2005.

- 42. Mrenna, S.: Event Generators.
<http://www.phys.psu.edu/~cteq/schools/summer04/mrenna/mrenna.pdf>.
- 43. Baer, H. et al.: Les Houches Guidebook to Monte Carlo Generators for Hadron Collider Physics. [hep-ph/0403045](#), 2004.
- 44. Sjostrand, T. et al.: PYTHIA 6.3 Physics and Manual. [hep-ph/0308153](#), 2003.
- 45. Altarelli, G. and Parisi, G.: Asymptotic freedom in parton language. Nucl. Phys. B, 126:298, 1977.
- 46. Gribov, V. and Lipatov, L.: . Sov. J. Nucl. Phys., 15:438, 1972.
- 47. Dokshitzer, Y.: . Sov. J. Phys. JETP, 46:641, 1977.
- 48. Mangano, M.: Higher-order QCD processes via LO matrix-element MC's.
<http://www.wlap.org/file-archive/atlas/20040622-umwlap002-02-mangano.pdf>.
- 49. Mangano, M.: Merging multijet matrix elements and shower evolution in hadronic collisions.
<http://mlm.home.cern.ch/mlm/talks/lund-alpgen.pdf>.
- 50. Catani, S. et al.: QCD matrix elements + parton showers. JHEP11(2001)063, 2001.
- 51. Michelangelo, L. et al.: A Generator for Hard Multiparton Processes in Hadronic Collisions. [hep-ph/0206293](#), 2002.
- 52. Mangano, M.: ALPGEN V1.3.
<http://mlm.home.cern.ch/mlm/alpgen/>.
- 53. Maltoni, F.: MadEvent: Automatic Event Generation with MADGRAPH. [hep-ph/0208156](#), 2002.
- 54. Mrenna, S.: Mrenna Matched Datasets.
http://cepa.fnal.gov/personal/mrenna/Matched_Dataset_Description.html.
- 55. Mrenna, S. and Richardson, P.: Matching matrix elements and parton showers with HERWIG and PYTHIA. JHEP05(2004)040, 2004.

56. Private Communication with Suyong Choi (suyong@fnal.gov) and <http://mcfm.fnal.gov/>.
57. Ellis, K.: A critical review of vector boson + jets Monte Carlos.
http://www-cdf.fnal.gov/Tevatron-Connection/QCD_Theory_Keith.pdf.
58. Ellis, K.: Method 2 at NLO. [hep-ph/0405276](http://arxiv.org/abs/hep-ph/0405276), 2004.
59. Fisyak, Y. and Womersley, J.: D0 GEANT Simulation of the Total Apparatus Response. [D0 Note 3191](#), 1997.
60. CERN: GEANT - Detector Description and Simulation Tool. 1993.
61. Choi, S. et al.: ATHENA Root Tuple Maker.
<http://www-d0.fnal.gov/~suyong/d0-private/athenaweb/athena.htm>.
62. Bassler, U. et al.: Technical Description of the T42 Algorithm for the Calorimeter Noise Suppression. [D0 Note 4146](#), 2003.
63. Stone, A.: Bad Runs.
<http://www-clued0.fnal.gov/~alstone/D0Work/Athena/badruns/badruns.html>.
64. Verzocchi, M.: Inclusive Single EM Triggers for W/Z Analyses.
<http://www-d0.fnal.gov/Run2Physics/wz/d0private/triggers/EMtriggers.html>.
65. Jain, S.: Scale and Over-smearing for MC Electron. [D0 Note 4402](#), 2004.
66. Top Physics Working Group: D0 Top Analyses and Data Sample for the Winter Conferences 2004. [D0 Note 4419](#), 2004.
67. Chapin, D. et al.: Measurement of the $Z \rightarrow e^+e^-$ and $W \rightarrow e^\pm\nu$ Production Cross Sections with $|\eta| < 1.05$. 2005.
http://www-d0.fnal.gov/~gardnerj/wzcross_v3.2.pdf.
68. Private Communication with Paul Telford (ptelford@fnal.gov).
69. CDF Collaboration: First Measurement of Inclusive W and Z Cross Sections from Run II of the Tevatron Collider. *Phys. Rev. Lett.*, 94:091803, 2005.
<http://www-cdf.fnal.gov/physics/preprints/index.html>.

70. Heinmiller, J. and Varelas, N.: Jet Reconstruction Efficiency. D0 Note 4837, 2005.
71. Private Communication with Mathieu Agelou (agelou@fnal.gov).
72. L2 Trigger Group: L2 Web Page. 2004.
<http://www.pa.msu.edu/hep/d0/l2/>.
73. UIC L2 Trigger Group: L2 Alpha Board Information. 2003.
http://www-d0online.fnal.gov/www/groups/trigger/l2/online/uic_alpha/.
74. Adams, M. et al.: Level-2 Calorimeter Preprocessor Technical Design Report. D0 Note 3651, 1999.
75. Edmunds, D. et al.: Technical Design Report for the L2 Global Processor. D0 Note 3402, 1998.
76. Linnemann, J.: The Standard L2 Crate. 1999.
http://www.pa.msu.edu/hep/d0/ftp/l2/overview/l2_standard_crate.pdf.
77. VITA: VME64 Extensions for Physics and other Application. 1997.
<http://galileo.phys.virginia.edu/~rjh2j/l2beta/vme/vita23d13a.pdf>.
78. Baden, D. et al.: D0 Level 2 Magic Bus. 2004.
http://galileo.phys.virginia.edu/~rjh2j/l2beta/beta_docs/d0_magicbus.pdf.
79. Hirosky, R.: The L2 Beta Project.
<http://galileo.phys.virginia.edu/~rjh2j/l2beta/home.html>.
80. University of Maryland D0 Level 2 Group: A Home page for the MagicBus Transceiver card.
<http://www-d0online.fnal.gov/www/groups/trigger/l2/cards/MBT/index.html>.
81. Linnemann, J.: FIC Card - Fiber Input Converter.
<http://www.pa.msu.edu/hep/d0/l2/fic.htm>.
82. Miller, S. et al.: Level 2 Processor Board Technical Reference Manual. 1998.
http://galileo.phys.virginia.edu/~rjh2j/l2beta/alpha_docs/alphaboard_technical.pdf.
83. DEC: Digital Semiconductor AlphaPC 164 Motherboard. 1997.
http://www-d0online.fnal.gov/www/groups/trigger/l2/online/uic_alpha/l2alpha/.

84. DEC: Digital Semiconducor 21164 Alpha Microprocessor Hardware Reference Manual. 1997.
http://www-d0online.fnal.gov/www/groups/trigger/l2/online/uic_alpha/l2alpha/.
85. Compaq: Alpha Architecture Handbook. 1998.
http://www-d0online.fnal.gov/www/groups/trigger/l2/online/uic_alpha/l2alpha/.
86. DEC: Digital Semiconducor 21172 Core Logic Chipset Technical Reference Manual. 1996.
http://www-d0online.fnal.gov/www/groups/trigger/l2/online/uic_alpha/l2alpha/.
87. Tundra: Universe II User Manual. 1998.
http://www-d0online.fnal.gov/www/groups/trigger/l2/online/uic_alpha/l2alpha/.

VITA

Marc Buehler was born in Rheinfelden, Germany on March 2, 1970. He grew up in Rheinfelden where he attended public school. In 1992 he earned a B.S. in Astronomy from the University of Basel, Switzerland. In 1997 he received a M.S. in Physics from the University of Heidelberg, Germany. He was accepted into the graduate physics program at the University of Illinois at Chicago, USA in 1997. Under the direction of Dr. Nikos Varelas, he received the degree of Doctor of Philosophy in Physics in 2005.